

# TEST ELEMENTS, RETRACTS AND AUTOMORPHIC ORBITS

SHENG-JUN GONG AND JIE-TAI YU

**Abstract.** Let  $A_2$  be a free associative or polynomial algebra of rank two over a field  $K$  of characteristic zero. Based on the degree estimate of Makar-Limanov and J.-T. Yu, we prove: 1) An element  $p \in A_2$  is a test element if  $p$  does not belong to any proper retract of  $A_2$ ; 2) Every endomorphism preserving the automorphic orbit of a nonconstant element of  $A_2$  is an automorphism.

## 1. INTRODUCTION AND MAIN RESULTS

In the sequel,  $K$  always denotes a field of characteristic zero, unless stated otherwise. Automorphisms (endomorphisms) always mean  $K$ -automorphisms ( $K$ -endomorphisms).

Let  $A_n$  be a free associative or polynomial algebra of rank  $n$  over  $K$ . An element  $p \in A_n$  is called a *test element* if every endomorphism of  $A_n$  fixing  $p$  is an automorphism. A subalgebra  $R$  of  $A_n$  is called a *retract* if there is an idempotent endomorphism  $\pi(\pi^2 = \pi)$  of  $A_n$  (called a *retraction* or a *projection*) such that  $\pi(A_n) = R$ . Test elements and retracts of the other algebras and groups are defined in a similar way. Test elements and retracts of algebras and groups have recently been studied in [3, 5, 6, 7, 12, 16, 20, 21, 22, 23, 24, 29, 30, 32, 33].

A test element does not belong to any proper retract for any algebra or group as the corresponding nonjective idempotent endomorphism is not an automorphism. The converse is proved by Turner [34] for free groups, by Mikhalev and Zolotykh [24] and by Mikhalev and J. -T. Yu [21, 22] for free Lie algebras and free Lie superalgebras respectively,

---

2000 *Mathematics Subject Classification.* Primary 16S10, 16W20; Secondary 13B10, 13F20.

*Key words and phrases.* Test element, retract, automorphic orbit, free associative algebra, degree estimate, polynomial algebra, coordinate.

Sheng-Jun Gong was partially supported by a University of Hong Kong Post-graduate Studentship.

Jie-Tai Yu was partially supported by an RGC-CERG Grant.

and by Mikhalev, Umirbaev and J. -T. Yu [19] for free nonassociative algebras. See also Mikhalev, Shpilrain and J. -T. Yu [16].

In view of the above, we may raise the following

**Conjecture 1.** If an element  $p \in A_n$  does not belong to any proper retract of  $A_n$ , then  $p$  is a test element.

Recently, V. Shpilrain and J. -T. Yu [33] proved Conjecture 1 for  $\mathbb{C}[x, y]$ . A key lemma in their proof is the degree estimate of Shestakov and Umirbaev [26], which plays a crucial role in the recent celebrated solution of the Nagata conjecture [27, 28] and the Strong Nagata conjecture [35].

More recently, Makar-Limanov and J. -T. Yu [18] developed a new combinatorial method based on the Lemma on radicals and obtained a sharp degree estimate for the ‘free’ case, namely, for a free associative algebra or a polynomial algebra over a field of characteristic zero. It has found applications for automorphisms and coordinates of polynomial and free associative algebras. See S.-J. Gong and J.-T. Yu [9].

Now we consider another related problem. In an algebra or a group, certainly an automorphism preserves the automorphic orbit of an element  $p$ . The converse is proved by Shpilrain [31] and Ivanov [10] for free groups of rank two, by D. Lee [14] for free groups of any rank, by Mikhalev and J. -T. Yu [22] for free Lie algebras and by Mikhalev, Umirbaev and J.-T. Yu [19] for free non-associative algebras, by van den Essen and Shpilrain [7] for  $A_2$  when  $p$  is a coordinate, by Jelonek [11] for polynomial algebras over  $\mathbb{C}$  when  $p$  is a coordinate. For the related linear coordinate preserving problem, see, for instance, S.-J. Gong and J.-T. Yu [8]. See also the book [16].

In view of the above, we may raise the following

**Conjecture 2.** Let  $p \in A_n - K$ . Then any endomorphism of  $A_n$  preserving the automorphic orbit of  $p$  must be an automorphism.

Conjecture 2 has recently been settled affirmatively by J.-T. Yu [36] based on Shpilrain and J.-T. Yu’s characterization of test elements of  $\mathbb{C}[x, y]$  in [33] and the main result in Drensky and J.-T. Yu [6].

In this paper, based on the recent degree estimate of Makar-Limanov and J.-T.Yu [18], the main ideals and techniques in Drensky and J.-T.Yu [6], Shpilrain and J.-T. Yu [32, 33], and J.-T.Yu [36], we prove both Conjecture 1 and Conjecture 2 for  $n = 2$ . Our main results are

**Theorem 1.1.** *If an element  $p \in A_2$  does not belong to any proper retract of  $A_2$ , then  $p$  is a test element of  $A_2$ .*

Theorem 1.1 was proved by Shpilrain and J.-T.Yu [33] for  $A_2 = \mathbb{C}[x, y]$ .

**Theorem 1.2.** *If an endomorphism  $\phi$  of  $A_2$  preserves the automorphic orbit of a nonconstant element  $p \in A_2$ , then  $\phi$  is an automorphism of  $A_2$ .*

Theorem 1.2 was proved by J.-T.Yu [36] for  $A_2 = \mathbb{C}[x, y]$ .

Crucial to the proofs of the above two theorems are the following two results, which have their own interests.

**Theorem 1.3.** *Let  $p \in A_2$  has outer rank two. Then any injective endomorphism  $\phi$  of  $A_2$  is an automorphism if  $\phi(p) = p$ .*

Theorem 1.3 may be viewed as an analogue of a result in Turner [34] for free groups. It was proved for  $A_2 = \mathbb{C}[x, y]$  in J.-T.Yu [36] based on a result in Shpilrain and J.-T.Yu [33].

**Theorem 1.4.** *An element  $p(x, y) \in A_2$  belongs to a proper retract of  $A_2$  if  $p(x, y)$  is fixed by a noninjective endomorphism  $\phi$  of  $A_2$ . Moreover, in this case there exists a positive integer  $m$  such that  $\phi^m$  is a retraction of  $A_2$ .*

Theorem 1.4 was proved for  $A_2 = \mathbb{C}[x, y]$  in Drensky and J.-T.Yu [6].

## 2. PROOFS

The following two lemmas are Theorem 1.1 and Proposition 1.2 in Makar-Limanov and J.-T.Yu [18].

**Lemma 2.1.** *Let  $A_n = K\langle x_1, \dots, x_n \rangle$  be a free associative algebra over a field  $K$  of characteristic zero,  $f, g \in A$  be algebraically independent,  $f^+$  and  $g^+$  are algebraically independent, or  $f^+$  and  $g^+$  are algebraically dependent and neither  $\deg(f) \mid \deg(g)$  nor  $\deg(g) \mid \deg(f)$ ,  $p \in K\langle x, y \rangle$ . Then*

$$\deg(p(f, g)) \geq \frac{\deg[f, g]}{\deg(fg)} w_{\deg(f), \deg(g)}(p).$$

Here  $\deg$  is the total degree,  $w_{\deg(f), \deg(g)}(p)$  is the weighted degree of  $p$  when the weight of the first variable is  $\deg(f)$  and the weight of the second variable is  $\deg(g)$ ,  $f^+$  and  $g^+$  are the highest homogeneous components of  $f$  and  $g$  respectively, and  $[f, g] = fg - gf$  is the commutator of  $f$  and  $g$ .

**Lemma 2.2.** *Let  $A_n = K[x_1, \dots, x_n]$  be a polynomial algebra over a field  $K$  of characteristic zero,  $f, g \in A$  be algebraically independent,  $p \in K[x, y]$ . Then*

$$\deg(p(f, g)) \geq w_{\deg(f), \deg(g)}(p) \left[ 1 - \frac{(\deg(f), \deg(g))(\deg(fg) - \deg(J(f, g)) - 2)}{\deg(f)\deg(g)} \right].$$

Here  $\deg$  is the total degree,  $w_{\deg(f), \deg(g)}(p)$  is the weighted degree of  $p$  when the weight of the first variable is  $\deg(f)$  and the weight of the second variable is  $\deg(g)$ ,  $(\deg(f), \deg(g))$  is the greatest common divisor of  $\deg(f)$  and  $\deg(g)$ ,  $\deg(J(f, g))$  is the largest degree of non-zero Jacobian determinants of  $f$  and  $g$  with respect to two of  $x_1, \dots, x_n$ .

The following characterization of a proper retract of  $A_2$  was obtained by Shpilrain and J.-T. Yu [32] based on a result of Costa [3].

**Lemma 2.3.** *Let  $R$  be a proper retract of  $A_2$ . Then  $R = K[r]$  for some  $r \in A_2$ . Moreover, there exists an automorphism  $\alpha$  of  $A_2$  such that  $\alpha(r) = x + w(x, y)$ , where  $w(x, y)$  belongs to the ideal of  $A_2$  generated by  $y$ .*

**Lemma 2.4.** *Let  $p \in A_2$  with outer rank 2 and  $f, g \in A_n$ . Then  $w_{\deg(f), \deg(g)}(p) \geq \deg(f) + \deg(g)$ .*

*Proof.* 1) If  $p$  contains a monomial containing both  $x$  and  $y$ , where  $i \neq 0, j \neq 0$ ,  $w_{\deg(f), \deg(g)}(p) \geq i(\deg(f)) + j(\deg(g)) \geq \deg(f) + \deg(g)$ .

2) Otherwise  $p$  must contain monomials  $x^i$  and  $y^j$  where  $i \geq 2, j \geq 2$ . Then  $w_{\deg(f), \deg(g)}(p) \geq 2 \max\{\deg(f), \deg(g)\} \geq \deg(f) + \deg(g)$ .  $\square$

**Lemma 2.5.** *Let  $A_n = K\langle x_1, \dots, x_n \rangle$  be a free associative algebra over an arbitrary field  $K$  of zero characteristic,  $f, g \in A$  be algebraically independent,  $p \in K\langle x, y \rangle$  has outer rank two. Then*

$$\deg(p(f, g)) \geq \deg[f, g].$$

*Proof.* Let 1) If  $f^+$  and  $g^+$  are algebraically independent; or  $f^+, g^+$  are algebraically dependent, but  $\deg(f) \nmid \deg(g)$  and  $\deg(g) \nmid \deg(f)$ . Then by Lemma 2.1 and Lemma 2.4,  $\deg(p(f, g)) \geq \deg[f, g]$ .

2) Otherwise there exists an automorphism  $\alpha$ , which is the composition of a sequence of elementary automorphisms, such that  $\alpha(f) = \bar{f}$ ,  $\alpha(g) = \bar{g}$ ,  $\bar{p} = \alpha^{-1}(p)$  satisfying the condition in 1). Then  $\deg(p(f, g)) = \deg(\bar{p}(\bar{f}, \bar{g})) \geq \deg[\bar{f}, \bar{g}] = \deg[f, g]$ .  $\square$

**Lemma 2.6.** *Let  $A_n = K[x_1, \dots, x_n]$  be a polynomial algebra over an arbitrary field  $K$  of zero characteristic,  $f, g \in A_n$  be algebraically independent,  $p \in K[x, y]$  has outer rank two. Then*

$$\deg(p(f, g)) \geq \deg(J(f, g)) + 2.$$

*Proof.* We may assume  $\deg(f) = m, \deg(g) = n$ . As  $p$  has outer rank 2, by Lemma 2.4 then  $p$  contains a monomial with both  $x$  and  $y$ , or contains monomials  $x^i$  and  $y^j$  where  $i \geq 2, j \geq 2$ .

1) Let  $f^+$  and  $g^+$  be algebraically independent.

a) If there exists a monomial in  $p$  containing both  $x$  and  $y$ , then  $\deg(p(f, g)) \geq \deg(f) + \deg(g) \geq \deg(J(f, g)) + 2$ ;

b) Otherwise  $p$  must have a monomial of  $x^i$  where  $i \geq 2$ , and another monomial  $y^j$  where  $j \geq 2$ , then  $\deg(p(f, g)) \geq 2 \max\{m, n\} \geq \deg(f) + \deg(g) \geq \deg(J(f, g)) + 2$ ;

2) Let  $f^+, g^+$  be algebraically dependent, and  $m \nmid n$  and  $n \nmid m$ .

c) If  $w_{\deg(f), \deg(g)}(p) < \text{lcm}(m, n)$ , then in  $p(f, g)$ ,  $f^+$  and  $g^+$  cannot cancel out, hence similar to the case 1 a),  $\deg(p(f, g)) \geq \deg(f) + \deg(g) \geq \deg(J(f, g)) + 2$ .

d) Otherwise  $w_{\deg(f), \deg(g)}(p) \geq \text{lcm}(m, n) = mn/(m, n)$ . We also have  $mn = (m, n)\text{lcm}(m, n) \geq (m, n)(m + n)$ . Hence  $\deg(p(f, g)) \geq \deg(J(f, g)) + 2$  by Lemma 2.2.

3) Let  $f^+, g^+$  be algebraically dependent, but  $m \mid n$  or  $n \mid m$ . Then by same process in the Proof 2) of Lemma 2.4, we may reduce to the above case 1) or case 2).  $\square$

**Lemma 2.7.** *Let  $\phi = (f, g)$  be an injective endomorphism of  $K\langle x, y \rangle$  but not an automorphism. Then  $\deg([\phi^k(x), \phi^k(y)]) \geq k + 2$  for  $k \geq 0$ .*

*Proof.*  $\deg[\phi^0(x), \phi^0(y)] = \deg[x, y] = 0 + 2$ . Since  $\phi$  is not an automorphism, by the well-known result of Dicks (see, Dicks [4], or Cohn [2]),  $\deg[\phi(x), \phi(y)] = \deg[f(x, y), g(x, y)] \geq \deg[x, y] + 1 = 1 + 2$ . Now the proof is concluded by

$$\deg[\phi^k(x), \phi^k(y)] =$$

$$\deg[f(\phi^{k-1}(x), \phi^{k-1}(y)), g(\phi^{k-1}(x), \phi^{k-1}(y))] \geq \deg[\phi^{k-1}(x), \phi^{k-1}(y)] + 1$$

(Note the above inequality follows by the aforementioned result of Dicks) and induction.  $\square$

**Lemma 2.8.** *Let  $\phi = (f, g)$  be an injective endomorphism of  $K[x, y]$  but not an automorphism and there exists an element  $p \in K[x, y]$  fixed by  $\phi$ . Then  $\deg(J(\phi^k(x), \phi^k(y))) \geq k$  for  $k \geq 0$ .*

*Proof.*  $\deg(J(\phi^0(x), \phi^0(y))) = \deg(J(x, y)) = 0$ . As  $\phi$  fixes  $p$ ,  $\phi$  is not an automorphism, by a result of Kraft [13] (see also Shpilrain and J.-T. Yu [32]),  $\deg(J(\phi(x), \phi(y))) = \deg(J(f, g)) \geq 1$ . By the chain rule for the Jacobian,

$$\begin{aligned} & \deg(J(\phi^k(x), \phi^k(y))) \\ &= \deg(J(f, g)(\phi^{k-1}(x), \phi^{k-1}(y))(J(\phi^{k-1}(x), \phi^{k-1}(y)))) \\ &\geq \deg(J(\phi^{k-1}(x), \phi^{k-1}(y))) + 1. \end{aligned}$$

The proof is concluded by induction.  $\square$

**Lemma 2.9.** *Let  $\phi = (f, g)$  be an injective endomorphism of  $A_2$  but not an automorphism. Then any element  $p \in A_2$  with outer rank 2 cannot be fixed by  $\phi$ .*

*Proof.* If  $p \in A_2$  with outer rank two fixed by  $\phi$ , then  $p(f, g) = p(\phi^k(x), \phi^k(y)) \geq k + 2$  for all  $k \geq 0$ . by Lemma 2.5 and Lemma 2.7 for noncommutative case; and by Lemma 2.6 and Lemma 2.8 for polynomial case. The contradiction completes the proof.  $\square$

### Proof of Theorem 1.3.

By Lemma 2.9.  $\square$

### Proof of Theorem 1.4.

The proof presented here is similar to the proof of the main Theorem in Drensky and J.-T. Yu [6].

Let  $p \in A_2 - \{0\}$  fixed by a noninjective endomorphism of  $A_2$ . Then  $\phi(x)$  and  $\phi(y)$  are algebraically dependent over  $K$ . Let us denote the image of  $\phi(A_2)$  by  $S = K[\phi(x), \phi(y)]$  (since  $\phi(x)$  and  $\phi(y)$  are algebraically dependent,  $\phi(x)$  and  $\phi(y)$  are in a polynomial algebra of rank one over  $K$  as a consequence of a result of Bergman [1] for noncommutative case and as a consequence of a result of Shestakov and Umirbaev [26] for polynomial case) and by  $Q(S)$  the field of fractions of  $S$ . Therefore the transcendence degree of  $Q(S)$  over  $K$  is 1. Let

$0 \neq q(x, y) \in (\text{Ker}(\phi)) \cap S$ . Since  $p(x, y)$  also belongs to  $S$ , the polynomials  $p$  and  $q$  are algebraically dependent and

$$h(p, q) = a_0(q)p^n + a_1(q)p^{n-1} + \dots + a_{n-1}(q)p + a_n(q) = 0$$

for an irreducible polynomial  $h(u, v) \in K[u, v]$  and  $a_i(t) \in K[t]$ ,  $i = 0, 1, \dots, n$ . Hence  $\phi(h(p, q)) = h(\phi(p), \phi(q)) = h(p, 0)$ ,

$$a_0(0)p^n + a_1(0)p^{n-1} + \dots + a_{n-1}(0)p + a_n(0) = 0.$$

Therefore  $a_0(0) = a_1(0) = \dots = a_n(0) = 0$ . Now the polynomials  $a_i(t)$  have no constant terms and  $h(u, v)$  is divisible by  $v$  which contradicts to the irreducibility of  $h(u, v)$ . Therefore  $(\text{Ker}(\phi)) \cap S = 0$  and  $\phi$  acts injectively on its image  $S$ . Hence we may extend the action of  $\phi$  on  $Q(S)$  (because  $a_1/b_1 = a_2/b_2$  in  $Q(S)$  is equivalent to  $a_1b_2 = a_2b_1$  and hence  $\phi(a_1/b_1) = \phi(a_1)/\phi(b_1) = \phi(a_2)/\phi(b_2) = \phi(a_2/b_2)$ ). By Lüroth's theorem (See, for instance, Schinzel [25]),  $Q(S) = K(w)$  for some  $w \in Q(S)$ . The automorphism  $\phi$  fixes  $p(x, y)$  and its extension  $\bar{\phi}$  on  $Q(S)$  fixes  $K(p)$ . Since  $w$  is algebraic over  $K(p)$ ,  $Q(S)$  is a finite dimensional vector space over  $K(p)$  and  $\bar{\phi}$  is a  $K(p)$ -linear operator of  $Q(S)$  with trivial kernel. Hence  $\bar{\phi}$  is invertible on  $Q(S)$  and we may consider  $\bar{\phi}$  as an automorphism of the finite field extension  $Q(S)$  over  $K(p)$  which fixes  $K(p)$ . By Galois theory ( $\bar{\phi}$  interchanges the roots of the minimal polynomial of  $w$  over  $K(p)$  and there are finite number of possibilities for  $\bar{\phi}(w)$ ),  $\bar{\phi}$  has finite order. Let  $\bar{\phi}^m = 1$ . Then  $\phi^{m+1}(r) = \phi^m(\phi(r)) = \bar{\phi}^m(\phi(r)) = \phi(r)$  for every  $r \in A_2$  and  $(\phi^m)^2 = \phi^{m+1}\phi^{m-1} = \phi\phi^{m-1} = \phi^m$ . Therefore  $\pi = \phi^m$  is a retraction (idempotent endomorphism) of  $A_2$  with a nontrivial kernel and  $\pi(p) = p$ . Hence  $p(x, y)$  is in the image of  $\pi$  which is a proper retract  $\pi(A_2)$  of  $A_2$ .  $\square$

### Proof of Theorem 1.1.

As  $p \in A_2$  does not belong to any proper retract of  $A_2$ , by Theorem 1.4, any endomorphism  $\phi$  of  $A_2$  fixing  $p$  must be injective. By Lemma 2.3, obviously  $p$  must have outer rank two, otherwise  $p$  would belong to a proper retract of  $A_2$ . By Theorem 1.3,  $\phi$  is an automorphism. Hence  $p$  is a test element of  $A_2$ .  $\square$

### Proof of Theorem 1.2.

The proof presented here is similar to the proof of the main result Theorem 1.4 in J.-T. Yu [36].

We may assume that  $\phi(p) = p$ . By the definition of the test element, we may assume  $p$  is not a test element. By Theorem 1.1, we may assume  $p$  belongs to a proper retract  $K[r]$  of  $A_2$ . By a result in J.-T. Yu [36], we may assume  $p$  has outer rank 2. By Theorem 1.3, we may assume  $\phi$  is non-injective. Suppose that  $p = f(r)$ , where  $f \in K[t] - K$ ,  $\deg(f) = m$ . By Theorem 1.4,  $\pi = \phi^m$  is a retraction of  $A_2$  to  $K[r]$ . As  $\phi$  preserves the automorphic orbit of  $p$ , so does  $\pi = \phi^m$ . Applying Lemma 2.3 (suppose  $\alpha(r) = x + w(x, y)$ , where  $w(x, y) \notin K[y]$  belongs to the ideal of  $A_2$  generated by  $y$ ,  $\alpha$  is some automorphism of  $A_2$ , replace  $r$  by  $\alpha(r)$ , and  $\pi$  by  $\alpha\pi\alpha^{-1}$ ), we have reduced our proof to the following

**Lemma 2.10.** *Let  $r = x + w(x, y)$ , where  $w(x, y)$  belongs to the ideal of  $A_2$  generated by  $y$  and  $w(x, y) \notin K[y]$ ,  $\pi$  the retraction of  $A_2$  onto  $K[r]$  defined by  $\pi(x) = x + w(x, y)$ ,  $\pi(y) = 0$ ,  $f \in K[t] - K$ . Then  $\pi$  does not preserve the automorphic orbit of  $f(r)$ .*

*Proof.* Suppose on the contrary,  $\pi$  preserves the automorphic orbit of  $f(r)$ . Then for any automorphism  $\alpha$  of  $A_2$ ,  $\pi\alpha(f(r)) = \beta(f(r)) \in K[r]$  for some automorphism  $\beta$  of  $A_2$ . Note that  $\pi\beta(f(r)) = \pi^2\beta(f(r)) = \pi\alpha(f(r)) = \beta(f(r))$ . By Theorem 1.4,  $\pi^{\deg(f)} = \pi$  is the retraction of  $A_2$  onto the retract  $K[\beta(r)]$  taking  $\beta(r)$  to  $\beta(r)$ . By hypothesis,  $\pi$  is also a retraction of  $A_2$  onto the retract  $K[r]$  taking  $r$  to  $r$ . This forces that  $\beta(r) = cr$  for some  $c \in K^*$ . We have concluded that for all automorphisms  $\alpha$  of  $A_2$ , there exists some  $c \in K^*$ , such that  $\pi\alpha(f(r)) = f(cr)$ .

Now we proceed the proof in two cases.

### 1. Noncommutative case: $A_2 = K\langle x, y \rangle$ .

Denote by  $\mathcal{C}$  the commutator ideal of  $K\langle x, y \rangle$ .

a) If  $w(x, y) \in \mathcal{C}$ , then take  $\alpha$  to be the automorphism of  $K\langle x, y \rangle$  defined by  $\alpha(x) = y + x^2$ ,  $\alpha(y) = x$ . Direct calculation shows that  $\pi\alpha(f(r)) = f(r^2 + w(r^2, r)) = f(r^2) \neq f(cr)$ , a contradiction.

b) If  $w(x, y) \notin \mathcal{C}$ , then  $w^a(x, y) = yv(x, y)$  for some  $v(x, y) \in K[x, y] - \{0\}$ . Here  $w^a(x, y) \in K[x, y]$  is the image of  $w(x, y)$  under the abelianization from  $K\langle x, y \rangle$  onto  $K[x, y]$ . Let  $M$  be a positive integer greater than  $\deg(v(x, y))$ , it is easy to see that  $x^M - y$  does not divide  $v(x, y)$  in  $K[x, y]$ . Let  $\alpha$  be the automorphism of  $K\langle x, y \rangle$  defined by  $\alpha(x) = x$ ,  $\alpha(y) = y + x^M$ . Then  $\pi\alpha(f(r)) = f(r + w(r, r^M)) =$

$f(r + r^M v(r, r^M))$ . As  $x^M - y$  does not divide  $v(x, y)$ ,  $v(r, r^M) \neq 0$ . Therefore  $\pi\alpha(f(r)) = f(r + r^M v(r, r^M)) \neq f(cr)$ , a contradiction.

## 2. Polynomial case: $A_2 = K[x, y]$ .

In this case we write  $w(x, y) = yq(x, y)$  where  $q(x, y) \notin K[y]$ . Let  $M$  be a positive integer greater than  $\deg(q(x, y))$ , it is easy to see that  $x^M - y$  does not divide  $q(x, y)$  in  $K[x, y]$ . Let  $\alpha$  be the automorphism of  $K[x, y]$  defined by  $\alpha(x) = x$ ,  $\alpha(y) = y + x^M$ . Then easy calculation shows that  $\pi\alpha(f(r)) = f(r + r^M q(r, r^M))$ . As  $x^M - y$  does not divide  $q(x, y)$ ,  $q(r, r^M) \neq 0$ . Therefore  $\pi\alpha(f(r)) = f(r + r^M q(r, r^M)) \neq f(cr)$ . The contradiction completes the proof.  $\square$

## REFERENCES

- [1] G. Bergman, *Centralizers in free associative algebras*, Trans. Amer. Math. Soc. **137** (1969), 327-344.
- [2] P. M. Cohn, *Free Rings and Their Relations*, 2nd Edition, London Mathematical Society Monograph, **19**, Academic Press, Inc. London, 1985.
- [3] D. Costa, *Retracts of polynomial rings*, J. Algebra **44** (1977), 492-502.
- [4] W. Dicks, *A commutator test for two elements to generate the free algebra of rank two*, Bull. London Math. Soc. **14** (1982), 48-51.
- [5] V. Drensky, J. -T. Yu, *Test Polynomials for automorphisms of polynomial and free associative algebras*, J. Algebra **207** (1998), 491-510.
- [6] V. Drensky, J. -T. Yu, *Retracts and test polynomials of polynomial algebras*, C. R. Acad. Bulgaria Sci. **55** (7) (2002), 11-14.
- [7] A. van den Essen, V. Shpilrain, *Some combinatorial questions about polynomial mappings*, J. Pure Appl. Algebra **119** (1997), 47-52.
- [8] S.-J. Gong, J.-T. Yu, *The linear coordinate preserving problem*, Preprint.
- [9] S.-J. Gong, J.-T. Yu, *The preimage of a coordinate*, Preprint.
- [10] S. Ivanov, *On endomorphisms of free groups that preserve primitivity*, Arch. Math. **72** (1999), 92-100.
- [11] Z. Jelonek, *A solution of the problem of van den Essen and Shpilrain*, J. Pure Appl. Algebra **137** (1999), 49-55.
- [12] Z. Jelonek, *Test polynomials*, J. Pure Appl. Algebra **147** (2000), 125-132.
- [13] H. Kraft, *On a question of Yosef Stein*, in 'Automorphisms of Affine Spaces (Curacao, 1994)' Proceedings of the Conference on Invertible Polynomial Maps held in Curacao, July 4-8, 1994. Ed. A. van den Essen, Kluwer Acad. Publ., Dordrecht, 1995, 225-229.
- [14] D. Lee, *Endomorphisms of free groups that preserve automorphic orbits*, J. Algebra **248** (2002), 230-236.
- [15] A. A. Mikhalev, V. Shpilrain, J. -T. Yu, *On endomorphisms of free algebras*, Algebra Colloq. **6** (1999), 241-248.
- [16] A. A. Mikhalev, V. Shpilrain, J. -T. Yu, *Combinatorial Methods: Free Groups, Polynomials, and Free Algebras*, CMS Books in Mathematics, Springer, New York, 2004.
- [17] L. Makar-Limanov, *On automorphisms of free algebra with two generators*, Funk. Analiz. Prilozh. **4** (1970), no. 3, 107-108. English translation: Funct. Anal. Appl. **4** (1970), 262-264.

- [18] L. Makar-Limanov, J. -T. Yu, *Degree estimate for two-generated subalgebras*, J. Euro. Math. Soc. (to appear)
- [19] A. A. Mikhalev, U. U. Umirbaev, J. -T. Yu, *Automorphic orbits of elements of free non-associative algebras*, J. Algebra **243** (2001), 198-223.
- [20] A. A. Mikhalev, U. U. Umirbaev, J. -T. Yu, *Generic, almost primitive and test elements of free Lie algebras*, Proc. Amer. Math. Soc. **130** (2002), 1303-1310.
- [21] A. A. Mikhalev, J. -T. Yu, *Test elements and retracts of free Lie algebras*, Commun. Algebra **25** (1997), 3283-3289.
- [22] A. A. Mikhalev, J. -T. Yu, *Test elements, retracts and automorphic orbits of free algebras*, Intern. J. Algebra Comput. **8** (1998), 295-310.
- [23] A. A. Mikhalev, J. -T. Yu, *Primitive, almost primitive, test, and  $\Delta$ -primitive elements of free algebras with the Nielsen-Schreier property*, J. Algebra **228** (2000), 603-623.
- [24] A. A. Mikhalev, A. A. Zolotykh, *Test elements for monomorphisms of free Lie algebras and Lie superalgebras*, Commun. Algebra **23** (1995), 4995-5001.
- [25] A. Schinzel, *Polynomials with Special Regard to Reducibility*, Encyclopedia of Mathematics, Cambridge University Press, Cambridge, 2000.
- [26] I. P. Shestakov, U. U. Umirbaev, *Poisson brackets and two-generated subalgebras of rings of polynomials*, J. Amer. Math. Soc. **17** (2004), 181-196.
- [27] I. P. Shestakov, U. U. Umirbaev, *The tame and the wild automorphisms of polynomial rings in three variables*, J. Amer. Math. Soc. **17** (2004), 197-227.
- [28] I. P. Shestakov, U. U. Umirbaev, *The Nagata Automorphism is Wild*, Proc. Nat. Acad. Sci. **100** (2003), No. 22, 12561-12563.
- [29] V. Shpilrain, *Recognizing automorphisms of the free groups*, Arch. Math. **62** (1994), 385-392.
- [30] V. Shpilrain, *Test elements for endomorphisms of free groups and algebras*, Israel. J. Math. **92** (1995), 307-316.
- [31] V. Shpilrain, *Generalized primitive elements of a free group*, Arch. Math. **71** (1998), 270-278.
- [32] V. Shpilrain, J. -T. Yu, *Polynomial retracts and the Jacobian conjecture*, Trans. Amer. Math. Soc. **352** (2000), 477-484.
- [33] V. Shpilrain, J. -T. Yu, *Test polynomials, retracts, and the Jacobian conjecture*, in Affine Algebraic Geometry, Contemp. Math. **369** (2005), 253-259, Amer. Math. Soc. Series, Providence, RI.
- [34] E. Turner, *Test words for automorphisms of free groups*, Bull. London Math. Soc. **28** (1996), 255-263.
- [35] U. U. Umirbaev, J. -T. Yu, *The Strong Nagata Conjecture*, Proc. Nat. Acad. Sci. **101** (2004), No. 13, 4352-4355.
- [36] J. -T. Yu, *Automorphic orbit problem for polynomial algebras*, J. Algebra (to appear)

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG, POKFULAM ROAD, HONG KONG SAR, CHINA  
*E-mail address:* `sjgong@hkusua.hku.hk`

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG, POKFULAM ROAD, HONG KONG SAR, CHINA  
*E-mail address:* `yujt@hkucc.hku.hk`, `yujietai@yahoo.com`