

# Packing Feedback Arc Sets in Tournaments Exactly

Xujin Chen<sup>a\*</sup> Guoli Ding<sup>b</sup> Wenan Zang<sup>c†</sup> Qiulan Zhao<sup>d‡</sup>

<sup>a</sup> Academy of Mathematics and Systems Science, Chinese Academy of Sciences  
Beijing 100190, China

<sup>b</sup> Mathematics Department, Louisiana State University  
Baton Rouge, LA 70803, USA

<sup>c</sup> Department of Mathematics, The University of Hong Kong  
Hong Kong, China

<sup>d</sup> Department of Mathematics, Nanjing University  
Nanjing 210093, China

## Abstract

Let  $T = (V, A)$  be a tournament with a nonnegative integral weight  $w(e)$  on each arc  $e$ . A subset  $F$  of arcs is called a *feedback arc set* (FAS) if  $T \setminus F$  contains no cycles (directed). A collection  $\mathcal{F}$  of FAS's (with repetition allowed) is called an *FAS packing* if each arc  $e$  is used at most  $w(e)$  times by the members of  $\mathcal{F}$ . The purpose of this paper is to give a characterization of all tournaments with the property that, for every nonnegative integral weight function  $w$  defined on  $A$ , the minimum total weight of a cycle is equal to the maximum size of an FAS packing.

**MSC 2000 subject classification.** Primary: 90C10, 90C27, 90C57.

**OR/MS subject classification.** Primary: Programming/graphs.

**Key words.** Tournament, feedback arc set, cycle, minimax relation, breadth-first search.

---

\*Supported in part by MOST of China under Grant 2018AAA0101002 and by Chinese Academy of Sciences under Grants XDA27000000 and ZDBS-LY-7008.

†Supported in part by the Research Grants Council of Hong Kong.

‡Corresponding author. E-mail: qiulanzhao@nju.edu.cn. Supported in part by NSF of China under Grant 11801266.

# 1 Introduction

Let  $G = (V, A)$  be a digraph with a nonnegative integral weight  $w(e)$  on each arc  $e$ . A subset  $F$  of arcs is called a *feedback arc set* (FAS) of  $G$  if  $G \setminus F$  contains no cycles (directed). The *FAS problem* is to find an FAS of  $G$  with minimum total weight, which can be naturally formulated as an integer program. One approach to this *NP*-hard problem is to consider its linear programming (LP) relaxation and explore integrality properties satisfied by its constraints. Let  $M$  be the cycle-arc incidence matrix of  $G$ , let  $\pi(G)$  denote the linear system  $Mx \geq \mathbf{1}$ ,  $x \geq \mathbf{0}$ , and let  $P$  denote the polyhedron defined by  $\pi(G)$ . We call  $P$  *integral* if it is the convex hull of all integral vectors contained in  $P$ . As is well known,  $P$  is integral iff the minimum in the LP-duality equation

$$\min\{w^T x : Mx \geq \mathbf{1}, x \geq \mathbf{0}\} = \max\{y^T \mathbf{1} : y^T M \leq w^T, y \geq \mathbf{0}\}$$

has an integral optimal solution, for every nonnegative integral vector  $w$  for which the optimum is finite. If, instead, the maximum in the equation satisfies this property, then the system  $\pi(G)$  is called *totally dual integral* (TDI). We say that  $G$  is *cycle ideal* (CI) if  $P$  is an integral polyhedron, and that  $G$  is *cycle Mengerian* (CM) if  $\pi(G)$  is a TDI system. As shown by Edmonds and Giles [15], total dual integrality implies primal integrality, so every CM digraph is CI and hence being CM can be more intuitively stated in terms of a minimax relation. A collection  $\mathcal{C}$  of cycles (with repetition allowed) is called a *cycle packing* of  $G$  if each arc  $e$  is used at most  $w(e)$  times by the members of  $\mathcal{C}$ . Let  $\nu_w(G)$  be the maximum size of a cycle packing, and let  $\tau_w(G)$  be the minimum total weight of an FAS. Then  $G$  is CM iff  $\nu_w(G) = \tau_w(G)$  for all nonnegative integral weight functions  $w$  defined on  $A$ . Note that a characterization of CI and CM digraphs can yield not only beautiful mathematical theorems but also a polynomial-time algorithm for the FAS problem on such digraphs, by a general theorem of Grötschel, Lovász, and Schrijver [17], so the study of these digraphs has both great theoretical interest and practical value. Despite tremendous research efforts, only some special classes of CI and CM digraphs [2, 3, 5, 7, 8, 18, 19, 23, 28] have been identified to date, and a complete characterization seems extremely hard to obtain.

The FAS problem remains *NP*-hard even when the input digraph  $G$  is a tournament; see Alon [1] and Charbit, Thomassé, and Yeo [10]. As this special version also arises in a rich variety of applications, it has been studied extensively from the combinatorial, statistical, and algorithmic points of view, and thus has produced a vast body of literature. Applegate, Cook, and McCormick [2] and Barahona, Fonlupt, and Mahjoub [3] independently proved that every tournament with five vertices is CM, thereby confirming a conjecture posed by both Barahona and Mahjoub [4] and Jünger [20]. We call a tournament *Möbius-free* if it contains none of  $K_{3,3}$ ,  $K'_{3,3}$ ,  $M_5$ , and  $M_5^*$  depicted in Figure 1 as a subgraph; these four Möbius ladders are actually the only obstructions to CI and CM tournaments.

**Theorem 1.1.** (Chen et al. [7, 8]) *For a tournament  $T$ , the following statements are equivalent:*

- (i)  $T$  is *Möbius-free*;
- (ii)  $T$  is *cycle ideal*; and
- (iii)  $T$  is *cycle Mengerian*.

Minimax relations in combinatorial optimization often appear in pairs (for example, a graph is perfect iff its complement is perfect). Given a minimax relation, a common practice in this




Figure 1. Forbidden Structures

field is to establish its blocker version. So a natural question to ask is: When does the minimax relation on packing and covering FAS's hold?

Let  $G = (V, A)$  and  $w$  be as given at the beginning of this section. We use  $N$  to denote the FAS-arc incidence matrix of  $G$ . A collection  $\mathcal{F}$  of FAS's (with repetition allowed) is called an *FAS packing* of  $G$  if each arc  $e$  is used at most  $w(e)$  times by the members of  $\mathcal{F}$ . Let  $\lambda_w(G)$  be the maximum size of an FAS packing, and let  $\mu_w(G)$  be the minimum total weight of a cycle (directed). Clearly,  $\lambda_w(G) \leq \mu_w(G)$ ; this inequality, however, need not hold with equality in general. We say that  $G$  is *FAS ideal* (FASI) if  $Nx \geq \mathbf{1}$ ,  $x \geq \mathbf{0}$  defines an integral polyhedron, and that  $G$  is *FAS Mengerian* (FASM) if  $Nx \geq \mathbf{1}$ ,  $x \geq \mathbf{0}$  is a TDI system. Again, by the aforementioned Edmonds-Giles theorem [15],  $G$  is FASM iff  $\lambda_w(G) = \mu_w(G)$  for every nonnegative integral weight function  $w$  defined on  $A$ . Since feedback arc sets are a type of combinatorial objects involving global structural properties, they are not so easily visualized as cycles and hence are more difficult to manipulate. Thus it is no surprise that packing FAS's in a digraph is harder than packing cycles.

The origin of FASM digraphs can be traced back to 1976, when Lucchesi and Younger [23] proved their min-max theorem on packing dicuts. We introduce some notions before proceeding. For each  $U \subseteq V$ , let  $\delta(U)$  denote the set of all arcs between  $U$  and  $V \setminus U$ , and let  $\delta^+(U)$  (resp.  $\delta^-(U)$ ) denote the set of arcs from  $U$  to  $V \setminus U$  (resp. from  $V \setminus U$  to  $U$ ) in  $G$ . A *dicut* is a set of arcs of the form  $\delta^+(U)$  for some subset  $U$  of  $V$  with  $\emptyset \neq U \neq V$  and with  $\delta^-(U) = \emptyset$ , which is also denoted by  $(U, V \setminus U)$ . A *dijoin* is a set of arcs that intersects every dicut. We can then define both *dicut packing* and *dijoin packing* in a similar way to cycle packing. The Lucchesi-Younger theorem [23] states that the maximum size of dicut packing is equal to the minimum total weight of a dijoin for all weight functions  $w$ . Edmonds and Giles [15] conjectured that the assertion remains true if we swap the terms dicut and dijoin; that is, the maximum size of dijoin packing is also equal to the minimum total weight of a dicut for all weight functions  $w$ . This conjecture

has been confirmed for several classes of digraphs such as source-sink connected digraphs [16, 26] and series-parallel digraphs [21]. The assertion of the general conjecture, however, was refuted by Schrijver [25]; more counterexamples have been found by Cornuéjols and Guenin [11] and Williams and Guenin [29]. Despite this, Woodall [30] strongly believed that the unweighted version of the Edmonds-Giles conjecture holds true. Motivated by this conjecture, Chudnovsky et al. [12] and Mészáros [24] have obtained several results on disjoint dijoins.

When restricted to a plane digraph, dicut and dijoin are dualized to cycle and feedback arc set, respectively. Thus the above Edmonds-Giles conjecture can be rephrased as saying that every planar digraph is FASM (a counterexample is the dual of Schrijver's digraph [25]), and Woodall's conjecture amounts to saying that the maximum number of disjoint feedback arc sets is equal to the length of a shortest cycle.

The purpose of this paper is to establish the blocker version of Theorem 1.1.

**Theorem 1.2.** *For a tournament  $T$ , the following statements are equivalent:*

- (i)  $T$  is Möbius-free;
- (ii)  $T$  is FAS ideal; and
- (iii)  $T$  is FAS Mengerian.

**Corollary 1.3.** *A tournament is cycle Mengerian iff it is FAS Mengerian iff it is Möbius-free.*

The reader is referred to [6] (resp. [9]) for a structural characterization of all undirected graphs (resp. tournaments) with the min-max relation on packing and covering feedback vertex sets and the corresponding blocker version [14, 13] (resp. [5]).

The remainder of this paper is organized as follows: In Section 2, we present a global structural description of Möbius-free strong tournaments. In Section 3, we establish the minimax relation on packing and covering FAS's in Möbius-free strong tournaments other than  $F_1$  and  $G_1$  (to be shown in Figures 4 and 5). In Section 4, we give a computer-assisted proof of the minimax relation on  $G_1$ , thereby completing the whole proof.

## 2 Global Structure

Our proof of Theorem 1.1 [7, 8] relies heavily on a structural description of Möbius-free strong tournaments, which continues to play an important role in the characterization of FAS Mengerian tournaments.

Let us recall some terminology and notations introduced in [7]. Let  $G = (V, A)$  be a digraph with a nonnegative integral weight  $w(e)$  on each arc  $e$ . For each  $v \in V$ , we use  $d_G^+(v)$  and  $d_G^-(v)$  to denote the out-degree and in-degree of  $v$ , respectively. We call  $v$  a *near-sink* of  $G$  if its out-degree is one, and call  $v$  a *near-source* if its in-degree is one. For simplicity, an arc  $e = (u, v)$  of  $G$  is also denoted by  $uv$ . Arc  $e$  is called *special* if  $u$  is a near-sink or  $v$  is a near-source of  $G$ . For each  $U \subseteq V$ , we use  $G[U]$  to denote the subgraph of  $G$  induced by  $U$ . Recall that  $G$  is called *weakly connected* if its underlying undirected graph is connected, and is called *strongly connected* or *strong* if each vertex is reachable from every other vertex. Clearly, a weakly connected digraph  $G$  is strong iff  $G$  has no dicut. A dicut  $(X, Y)$  is called *trivial* if  $|X| = 1$  or  $|Y| = 1$ . Furthermore,

a weakly connected digraph  $G$  is called *internally strong* if every dicut of  $G$  is trivial, and is called *internally 2-strong (i2s)* if  $G$  is strong and  $G \setminus v$  is internally strong for every vertex  $v$ .

Let  $T_i = (V_i, A_i)$  be a tournament, with  $|V_i| \geq 3$  for  $i = 1, 2$ . We say that  $T_1$  is *smaller* than  $T_2$  if  $|V_1| < |V_2|$ . Suppose that  $(a_1, b_1)$  is a special arc of  $T_1$  with  $d_{T_1}^+(a_1) = 1$  and  $(b_2, a_2)$  is a special arc of  $T_2$  with  $d_{T_2}^-(a_2) = 1$ . The *1-sum* of  $T_1$  and  $T_2$  over  $(a_1, b_1)$  and  $(b_2, a_2)$  is the tournament arising from the disjoint union of  $T_1 \setminus a_1$  and  $T_2 \setminus a_2$  by identifying  $b_1$  with  $b_2$  (the resulting vertex is denoted by  $b$ ) and adding all arcs from  $T_1 \setminus \{a_1, b_1\}$  to  $T_2 \setminus \{a_2, b_2\}$ . We call  $b$  the *hub* of the 1-sum. See Figure 2 for an illustration. Note that if  $|V_i| = 3$  for  $i = 1$  or  $2$ , then  $T_i$  is a triangle (a directed cycle of length three), and thus  $T = T_{3-i}$ .




Figure 2. 1-sum of  $T_1$  and  $T_2$ .

In our original definition of 1-sum [7, 8], we assume that  $T_i = (V_i, A_i)$  is strong for  $i = 1, 2$ ; this assumption is removed here just for more convenience. The lemma below asserts that these two definitions are equivalent when restricted to a strong tournament  $T$ .

**Lemma 2.1.** *Suppose a strong tournament  $T$  is the 1-sum of two tournaments  $T_1$  and  $T_2$ . Then the following statements hold:*

- (i) *Both  $T_1$  and  $T_2$  are strong; and*
- (ii) *Both  $T_1$  and  $T_2$  are subtournaments of  $T$ .*

As the proof is completely straightforward, we omit it here. Lemma 2.1 allows us to assume that  $a_i$  is a vertex of  $T_{3-i} \setminus a_{3-i}$  for  $i = 1, 2$  in the definition of the 1-sum.

The following lemma (see Lemma 2.2 in [7]) states that being Möbius-free is closed under taking 1-sums.

**Lemma 2.2.** *Suppose a strong tournament  $T$  is a 1-sum of two tournaments  $T_1$  and  $T_2$ . Then  $T$  is Möbius-free iff both  $T_1$  and  $T_2$  are Möbius-free.*

Let  $C_3$  (resp.  $F_0$ ) denote the strong tournament with three (resp. four) vertices (see Figure 3), let  $F_1, F_2, F_3, F_4, F_5$  be the five tournaments depicted in Figure 4, and let  $G_1, G_2, G_3$  be the three tournaments shown in Figure 5. We reserve the symbols

$$\mathcal{T}_0 = \{C_3, F_0, F_1, F_2, F_3, F_4, G_1, G_2, G_3\}$$

and

$$\mathcal{T}_1 = \{C_3, F_0, F_2, F_3, F_4, G_2, G_3\} = \mathcal{T}_0 \setminus \{F_1, G_1\}.$$

In [7] we have obtained the following structural descriptions of Möbius-free tournaments.




Figure 3. Strong tournaments with three or four vertices.




Figure 4.  $v_1v_2, v_5v_1 \in F_1$ ;  $v_2v_1, v_1v_5 \in F_2$ ;  $v_2v_1, v_5v_1 \in F_3$ ;  $v_6v_2 \in F_4$ ;  $v_2v_6 \in F_5$ .




Figure 5.  $v_6v_4 \in G_2$  and  $v_4v_6 \in G_3$ .




Figure 6. A minimal tournament involved in 1-sum

**Theorem 2.3.** (Chen et al. [7]) *Let  $T = (V, A)$  be an i2s tournament with  $|V| \geq 3$ . Then  $T$  is Möbius-free iff  $T \in \mathcal{T}_0$ .*

**Theorem 2.4.** (Chen et al. [7]) *Let  $T = (V, A)$  be a Möbius-free strong tournament with  $|V| \geq 3$ . Then either  $T \in \{F_1, G_1\}$  or  $T$  can be obtained by repeatedly taking 1-sums starting from the tournaments in  $\mathcal{T}_1$ .*

Let  $F_6$  be the tournament depicted in Figure 6 and let

$$\mathcal{T}_2 = \{F_0, F_2, F_3, F_4, F_6, G_2, G_3\}.$$

Then  $\mathcal{T}_2 = (\mathcal{T}_1 \setminus \{C_3\}) \cup \{F_6\}$ . The following is a simplified version of Lemma 2.4 in [8], which follows instantly from Lemma 2.1.


**Lemma 2.5.** *Let  $T = (V, A)$  be a Möbius-free strong tournament. Suppose  $T$  is a 1-sum of two smaller tournaments  $T_1$  and  $T_2$  such that  $T_2$  is minimal (with respect to vertex set inclusion). Then  $T_2 \in \mathcal{T}_2$ .*

Note that neither  $F_1$  nor  $G_1$  has a near-sink while every other tournament in  $\mathcal{T}_0$  does. Thus the above three results imply the following.

**Corollary 2.6.** *Let  $T = (V, A)$  be a Möbius-free strong tournament, with  $T \notin \{C_3, F_1, G_1\}$ . Then  $T$  can be constructed from a tournament in  $\{F_0, F_2, F_3, F_4, G_2, G_3\}$  by repeatedly 1-summing a tournament in  $\mathcal{T}_2$ .*

So far we have exhibited some local structural properties satisfied by Möbius-free strong tournaments. Due to the global nature of feedback arc sets, we need a description of entire tournaments in order to establish the desired minimax relation.

**Theorem 2.7.** *A strong tournament  $T = (V, A)$  other than  $F_1$  and  $G_1$  is Möbius-free iff it satisfies the following description:*



and all other arcs are directed from “left” to “right”.

Figure 7. Global structure

Let  $\mathcal{P}$  denote the class of all strong tournaments  $T$  described in the above theorem. We call  $A_1, A_2, \dots, A_m$  *vertical blocks* of  $T$ , call  $B_1, B_2, \dots, B_{m-1}$  *horizontal blocks* of  $T$ , and call  $v_1, v_2, \dots, v_m$  the *join vertices* of  $T$ . Let  $\mathcal{Q}$  consist of all tournaments  $G$  whose vertex set can be partitioned into  $U_0, U_1, \dots, U_k$ , such that  $|U_0| = 1$ ,  $G[U_i]$  is either a singleton or a triangle for  $1 \leq i \leq k$ , and the arcs between  $U_i$  and  $U_j$  are all directed to  $U_j$  for  $1 \leq i < j \leq k$ . Let  $v$  be the vertex in  $U_0$ . We call  $v$  the *center* of  $G$  and call  $G[U_i \cup \{v\}]$  a *building block* of  $G$  centered at  $v$  for  $1 \leq i \leq k$ . Clearly, each vertical block  $A_i$  of  $T$  belongs to  $\mathcal{Q}$ . We reserve the symbols  $A_{i,1}, A_{i,1}, \dots, A_{i,n_i}$  for the building blocks of  $A_i$  centered at  $v_i$ , where  $n_i \geq 0$ .

Let us prove three technical lemmas before presenting a proof of Theorem 2.7.

**Lemma 2.8.** *Every tournament in  $\{C_3\} \cup \mathcal{T}_2$  belongs to  $\mathcal{P}$ .*

**Proof.** The statement holds trivially for  $C_3$ . As shown in Figure 8 (where the missing arcs are all directed from left to right),  $F_0$  can be expressed in two ways, with  $m = 2$  and  $m = 1$ , respectively;  $F_3$  and  $F_4$  can be expressed with  $m = 2$ , while  $F_2, F_6, G_2$  and  $G_3$  can be expressed with  $m = 1$ . ■




Figure 8. Tournaments in  $\mathcal{T}_2$

**Lemma 2.9.** *Let  $G$  be a strong tournament on five vertices with a near-sink or a near-source. Then  $G$  is  $F_2$  or  $F_3$  or a 1-sum of two copies of  $F_0$ .*

**Proof.** If  $G$  is *i2s*, then  $G \in \{F_1, F_2, F_3\}$  by Theorem 2.3 and hence  $G$  is  $F_2$  or  $F_3$ , because  $F_1$  contains no near-sink nor near-source. So we assume that  $G$  is not *i2s*. By definition,  $G \setminus v$  has a dicut  $(X, Y)$  with  $|X| = |Y| = 2$  for some vertex  $v$ . Since  $G$  is strong, there exist a vertex  $a_1$  in  $Y$  and a vertex  $a_2$  in  $X$ , such that both  $(a_1, v)$  and  $(v, a_2)$  are arcs of  $G$ . Let  $T_1$  be the subtournament of  $G$  induced by the vertex subset  $X \cup \{a_1, v\}$  and let  $T_2$  be the subtournament of  $G$  induced by the vertex subset  $Y \cup \{a_2, v\}$ . Then  $T_1$  and  $T_2$  are two copies of  $F_0$  and  $G$  is the 1-sum of  $T_1$  and  $T_2$  over arcs  $(a_1, v)$  and  $(v, a_2)$ . It is routine to check that each of these 1-sums (three in total) contains a near-sink or a near-source. ■

In what follows,  $\mathcal{R}_5$  is the set of all strong tournaments on five vertices with a near-sink or a near-source, and  $F_6^*$  arises from  $F_6$  by reversing the direction of each arc.

**Lemma 2.10.** *Let  $G = (V, A)$  be a strong tournament in  $\mathcal{Q}$  with at least three vertices. Then either  $G \in \{C_3, F_0\}$  or  $G$  can be obtained by repeatedly taking 1-sums starting from tournaments in  $\{F_0, F_6, F_6^*, G_2, G_3\} \cup \mathcal{R}_5$ , such that the hubs of these sums are always the center of  $G$ .*

**Proof.** Let  $v$  be the center of  $G$  and let  $H_1, H_2, \dots, H_k$  be the building blocks of  $G$  centered at  $v$ , where the arcs between  $H_i \setminus v$  and  $H_j \setminus v$  are all directed to  $H_j$  for  $1 \leq i < j \leq k$ . We proceed by induction on  $k$ . If  $k = 1$ , then  $G = F_0$  and hence the statement holds trivially. So we assume that  $k \geq 2$  and set  $X_i := V(H_i)$  for  $1 \leq i \leq k$ .

We first assume that  $|H_1| = 4$ . Let  $a_1$  be an in-neighbor of  $v$  in  $G \setminus X_1$  and  $a_2$  be an out-neighbor of  $v$  in  $H_1$  (such  $a_1$  and  $a_2$  exist, as  $G$  is strong). Let  $T_1$  and  $T_2$  be the subtournaments of  $G$  induced by  $X_1 \cup \{a_1\}$  and  $(V \setminus X_1) \cup \{v, a_2\}$ , respectively. Then  $G$  is the 1-sum of  $T_1$  and  $T_2$  over arcs  $(a_1, v)$  and  $(v, a_2)$ . Note that  $T_1 \in \mathcal{R}_5$  and  $T_2 \in \mathcal{Q}$ . By induction hypothesis, either  $T_2 \in \{C_3, F_0\}$  or  $T_2$  can be obtained by repeatedly taking 1-sums starting from tournaments in  $\{F_0, F_6, F_6^*, G_2, G_3\} \cup \mathcal{R}_5$ , such that the hubs of these sums are always  $v$ . Clearly,  $G = T_1$  when  $T_2 = C_3$ . Therefore  $G$  can be obtained by repeatedly taking 1-sums starting from tournaments in  $\{F_0, F_6, F_6^*, G_2, G_3\} \cup \mathcal{R}_5$ , such that the hubs of these sums are always the center of  $G$ .

Next, we assume that  $|H_1| = 2$ . If  $k = 2$ , then  $G$  is either a  $C_3$  or a strong tournament on five vertices with a near-source, so the desired statement holds trivially. Thus we may further assume that  $k \geq 3$ . Let  $a_1$  be an in-neighbor of  $v$  in  $G \setminus (X_1 \cup X_2)$  and  $a_2$  be an out-neighbor of  $v$  in  $H_1 \cup H_2$  (such  $a_1$  and  $a_2$  exist, as  $G$  is strong). Let  $T_1$  and  $T_2$  be the tournaments induced by  $X_1 \cup X_2 \cup \{a_1\}$  and  $(V \setminus (X_1 \cup X_2)) \cup \{v, a_2\}$ , respectively. From Figure 8 we see that  $T_1 \in \{F_0, F_6, F_6^*, G_2, G_3\}$ . (Note that  $T_1 = F_6^*$  if  $|H_2| = 4$  and  $v$  is a source of  $H_2$ .) Clearly,  $T_2 \in \mathcal{Q}$  and  $G$  is the 1-sum of  $T_1$  and  $T_2$  over arcs  $(a_1, v)$  and  $(v, a_2)$ . By the induction hypothesis, either  $T_2 \in \{C_3, F_0\}$  or  $T_2$  can be obtained by repeatedly taking 1-sums starting from tournaments in  $\{F_0, F_6, F_6^*, G_2, G_3\} \cup \mathcal{R}_5$ , such that the hubs of these sums are always  $v$ . Clearly,  $G = T_1$  when  $T_2 = C_3$ . Therefore  $G$  can be obtained by repeatedly taking 1-sums starting from tournaments in  $\{F_0, F_6, F_6^*, G_2, G_3\} \cup \mathcal{R}_5$ , such that the hubs of these sums are always the center of  $G$ . ■

Now we are ready to establish the main result of this section.

**Proof of Theorem 2.7.** Let us first show the “if” part. Let  $T = (V, A)$  be a tournament in  $\mathcal{P}$  as described in Figure 7, with vertical blocks  $A_1, A_2, \dots, A_m$ , horizontal blocks  $B_1, B_2, \dots, B_{m-1}$ , and join vertices  $v_1, v_2, \dots, v_m$ ; subject to this, we assume that  $m$  is minimum (the choices of  $A_i$  and  $B_i$  may not be unique). This assumption implies that  $|A_1| \geq 2$  and  $|A_m| \geq 2$ . Furthermore,

(1)  $|A_1| \geq 3$  if  $|B_1| \leq 3$ , for otherwise, let  $A'_2$  be the subtournament of  $T$  induced by all vertices in  $A_1 \cup A_2 \cup B_1$ . Then  $T$  can be depicted as in Figure 7, with vertical blocks  $A'_2, A_3, \dots, A_m$  and horizontal blocks  $B_2, B_3, \dots, B_{m-1}$ , contradicting the minimality assumption on  $m$ .

By Theorem 2.3,  $C_3, F_0, F_2, G_2$  and  $G_3$  are all Möbius-free. Note that  $F_6$  is the 1-sum of  $F_2$  and  $F_0$  (see Figure 6). By Lemma 2.2,  $F_6$  is also Möbius-free and hence so is its reverse  $F_6^*$ . Since  $A_i \in \mathcal{Q}$ , it follows from Lemma 2.2 and Lemma 2.10 that

(2)  $A_i$  is Möbius-free for  $1 \leq i \leq m$ .

We propose to show, by induction on  $m+n$ , that  $T$  is Möbius-free, where  $n = |V|$ . If  $m = 1$ , then  $T = A_1$ , so the statement is true by (2). If  $n \leq 5$ , trivially the statement holds. Thus we may assume that  $m \geq 2$  and  $n \geq 6$ .

Consider the case when  $|A_1| = 2$ . Now  $|B_1| = 4$  by (1). Besides, there are at least two vertices outside  $A_1 \cup B_1$ , for otherwise,  $T = F_4$  (see Figure 8), which is Möbius-free by Theorem 2.3. Let  $a_1$  be an in-neighbor of  $v_2$  outside  $A_1 \cup B_1$  and let  $a_2$  be an out-neighbor of  $v_2$  in  $B_1$ .

Let  $T_1$  be the subtournament of  $T$  induced by all vertices in  $V(A_1 \cup B_1) \cup \{a_1\}$  and let  $T_2$  be the subtournament of  $T$  induced by all vertices outside  $V(A_1 \cup B_1) \setminus \{v_2, a_2\}$ . Then  $T_1$  is  $F_4$  (see Figure 8),  $T_2$  is a tournament in  $\mathcal{P}$  with  $m - 1$  vertical blocks, and  $T$  is the 1-sum of  $T_1$  and  $T_2$  over arcs  $(a_1, v_2)$  and  $(v_2, a_2)$ . By induction hypothesis,  $T_2$  is Möbius-free and hence so is  $T$  by Lemma 2.2.

It remains to consider the case when  $|A_1| \geq 3$ . Let  $a_1$  be an in-neighbor of  $v_1$  outside  $A_1$  and let  $a_2$  be an out-neighbor of  $v_1$  in  $A_1$  (such  $a_1$  and  $a_2$  exist, as  $T$  is strong). Let  $T_1$  be the subtournament of  $T$  induced by all vertices in  $V(A_1) \cup \{a_1\}$  and let  $T_2$  be the subtournament of  $T$  induced by all vertices outside  $A_1 \setminus \{a_2, v_1\}$ . Note that  $T_i \in \mathcal{P}$ ,  $4 \leq |T_i| < n$  for  $i = 1, 2$ , and  $T$  is the 1-sum of  $T_1$  and  $T_2$  over arcs  $(a_1, v_1)$  and  $(v_1, a_2)$ . By induction hypothesis,  $T_i$  is Möbius-free for  $i = 1, 2$  and hence so is  $T$  by Lemma 2.2. This establishes the “if” part.

Let us now proceed to the “only if” part. Let  $T = (V, A)$  be a strong Möbius-free tournament other than  $F_1$  and  $G_1$ . We aim to show, by induction on  $n = |V|$ , that  $T \in \mathcal{P}$ . If  $T$  is  $i2s$ , then  $T \in \{C_3\} \cup \mathcal{T}_2$  by Theorem 2.3 and hence  $T \in \mathcal{P}$  by Lemma 2.8. So we assume that  $T$  is not  $i2s$ . Then  $T$  a 1-sum of two smaller tournaments  $T_1$  and  $T_2$  over two special arcs  $(a_1, b_1)$  and  $(b_2, a_2)$ , such that  $T_2 \in \mathcal{T}_2$  by Lemma 2.5. Recall that  $a_i$  and  $b_i$  are vertices of  $T_i$  for  $i = 1, 2$ .

By induction hypothesis,  $T_1$  is as described in Figure 7, with vertical blocks  $A_1, A_2, \dots, A_m$ , horizontal blocks  $B_1, B_2, \dots, B_{m-1}$ , and join vertices  $v_1, v_2, \dots, v_m$ ; subject to this, we assume that  $m$  is minimum. This assumption implies that

(3)  $|A_1| \geq 2$  and  $|A_m| \geq 2$ .

If  $m = 1$ , then  $T_1 = A_1 \in \mathcal{Q}$ . Since  $a_2$  is a near-source of  $T_2$ ,  $(b_2, a_2)$  is the left-most arc of the corresponding tournament shown in Figure 8. Thus the 1-sum of  $T_1$  and a tournament in  $\{F_0, F_2, F_6, G_2, G_3\}$  belongs to  $\mathcal{Q}$ , and the 1-sum of  $T_1$  and a tournament in  $\{F_3, F_4\}$  belongs to  $\mathcal{P}$  with two vertical blocks. Hence  $T \in \mathcal{P}$ , as desired. So we assume that  $m \geq 2$ . Since  $a_1$  is a near-sink of  $T_1$ , it belongs to  $B_{m-1} \cup A_m$ . If  $a_1 \in V(B_{m-1} \setminus v_m)$ , then  $|B_{m-1}| = 2$  or 3 and  $V(A_m) = \{v_m\}$ , contradicting (3). If  $a_1 = v_m$ , then  $B_{m-1}$  consists of only one arc  $(v_m, v_{m-1}) = (a_1, b_1)$  and  $v_m$  is a sink of  $A_m$ . Thus we can combine  $A_{m-1}$ ,  $B_{m-1}$  and  $A_m$  to form a new  $A'_{m-1}$  and depict  $T_1$  as in Figure 7, with vertical blocks  $A_1, A_2, \dots, A_{m-2}, A'_{m-1}$  and horizontal blocks  $B_1, B_2, \dots, B_{m-2}$ , contradicting the minimality assumption on  $m$ . So  $a_1 \in V(A_m \setminus v_m)$ . Let  $A_{m,1}, A_{m,2}, \dots, A_{m,n_m}$  be the building blocks of  $A_m$  centered at  $v_m$ . Again, since  $a_1$  is a near-sink of  $T_1$ , we obtain

(4)  $a_1$  is contained in  $(A_{m,n_m-1} \cup A_{m,n_m}) \setminus v_m$ .

For simplicity, in the remainder of this proof, we frequently define  $B_m, A_{m+1}$ , etc. in terms of vertex sets only. For example, by  $B_m = \{b_1, v_m\}$  we mean that  $B_m$  is the tournament with vertex set  $\{b_1, v_m\}$ .

We first assume that  $a_1 \in V(A_{m,n_m-1})$ . Then  $A_{m,n_m-1}$  consists of only one arc  $(v_m, a_1)$  and  $A_{m,n_m}$  consists of only one arc  $(b_1, v_m)$  (as  $T_1$  is strong by Lemma 2.1). If  $T_2 \neq F_4$  (possibly  $T_2 = F_3$ ; see Figure 8), then  $T \in \mathcal{P}$  with the join vertices  $v_1, \dots, v_m, v_{m+1} := b_1$  and with new blocks  $A_m := A_m \setminus \{a_1, b_1\}$ ,  $B_m = \{b_1, v_m\}$ , and  $A_{m+1} = T_2 \setminus a_2$ ; if  $T_2 = F_4$  (see Figure 8), then  $T$  is in  $\mathcal{P}$  with join vertices  $v_1, \dots, v_m, v_{m+1} := b_1, v_{m+2} := z_3$  and with new blocks  $A_m := A_m \setminus \{a_1, b_1\}$ ,  $B_m = \{b_1, v_m\}$ ,  $A_{m+1} = \{b_1\}$ ,  $B_{m+1} = \{b_1, z_1, z_3, z_4\}$ , and  $A_{m+2} = \{z_2, z_3\}$ .

Next, we assume that  $a_1 \in V(A_{m,n_m})$ . Consider the case when  $|A_{m,n_m}| = 2$ . Now  $A_{m,n_m}$  consists of arc  $(a_1, v_m)$  only and  $b_1 = v_m$ . If  $T_2 \in \{F_0, F_2, F_6, G_2, G_3\}$ , where  $F_0$  corresponds to  $m = 1$  in Figure 8, then  $T \in \mathcal{P}$  with join vertices  $v_1, v_2, \dots, v_m$  and with new block  $A_m$

equal to the subtournament of  $T$  induced by all vertices in  $(A_m \setminus a_1) \cup (T_2 \setminus \{a_2, b_2\})$ . If  $T_2 = F_0$  corresponds to  $m = 2$  in Figure 8, then  $T \in \mathcal{P}$  with join vertices  $v_1, \dots, v_m, v_{m+1} := z_4$  and with new blocks  $B_m = \{z_4, z_1\}$ ,  $A_m := A_m \setminus a_1$ , and  $A_{m+1} = \{z_3, z_4\}$ . If  $T_2 = F_3$ , then  $T \in \mathcal{P}$  with join vertices  $v_1, \dots, v_m, v_{m+1} := z_3$  and with new blocks  $B_m = \{z_3, z_4, z_5\}$ ,  $A_m := A_m \setminus a_1$ , and  $A_{m+1} = \{z_1, z_3\}$ . If  $T_2 = F_4$ , then  $T \in \mathcal{P}$  with join vertices  $v_1, \dots, v_m, v_{m+1} := z_3$  and with new blocks  $B_m = \{z_1, z_3, z_4, z_5\}$ ,  $A_m := A_m \setminus a_1$ , and  $A_{m+1} = \{z_2, z_3\}$ .

It remains to consider the case when  $|A_{m,n_m}| = 4$ . Now  $A_{m,n_m} \setminus v_m$  is a triangle  $a_1 b_1 c_1 a_1$ . Since  $a_1$  is a near-sink,  $(v_m, a_1)$  is an arc of  $T_1$ . Since  $T_1$  is strong, at least one of the two arcs between  $v_m$  and  $\{b_1, c_1\}$  is directed to  $v_m$ .

Suppose  $(b_1, v_m)$  and  $(c_1, v_m)$  are two arcs of  $T_1$ . If  $T_2 \neq F_4$  (possibly  $T_2 = F_3$ ; see Figure 8), then  $T \in \mathcal{P}$  with join vertices  $v_1, \dots, v_m, v_{m+1} := b_1$  and with new blocks  $A_m := A_m \setminus \{a_1, b_1, c_1\}$ ,  $B_m = \{b_1, c_1, v_m\}$ , and  $A_{m+1} = T_2 \setminus a_2$ . If  $T_2 = F_4$  (see Figure 8), then  $T \in \mathcal{P}$  with join vertices  $v_1, \dots, v_m, v_{m+1} := b_1, v_{m+2} := z_3$  and with new blocks  $A_m := A_m \setminus \{a_1, b_1, c_1\}$ ,  $B_m = \{b_1, c_1, v_m\}$ ,  $A_{m+1} = \{b_1\}$ ,  $B_{m+1} = \{b_1, z_1, z_3, z_4\}$ , and  $A_{m+2} = \{z_2, z_3\}$ .

So we assume that exactly one of the two arcs between  $v_m$  and  $\{b_1, c_1\}$  is directed to  $v_m$ . When  $(b_1, v_m)$  and  $(v_m, c_1)$  are two arcs of  $T_1$ , we see that if  $T_2 \neq F_4$  (possibly  $T_2 = F_3$ ; see Figure 8), then  $T \in \mathcal{P}$  with join vertices  $v_1, \dots, v_m, v_{m+1} := b_1$  and with new blocks  $A_m := A_m \setminus \{a_1, b_1, c_1\}$ ,  $B_m = \{b_1, v_m\}$ , and  $A_{m+1}$  equal to the subtournament of  $T$  induced by  $\{c_1\} \cup V(T_2 \setminus a_2)$ ; if  $T_2 = F_4$  (see Figure 8), then  $T \in \mathcal{P}$  with join vertices  $v_1, \dots, v_m, v_{m+1} := b_1, v_{m+2} := z_3$  and with new blocks  $A_m := A_m \setminus \{a_1, b_1, c_1\}$ ,  $B_m = \{b_1, v_m\}$ ,  $A_{m+1} = \{b_1, c_1\}$ ,  $B_{m+1} = \{b_1, z_1, z_3, z_4\}$ , and  $A_{m+2} = \{z_2, z_3\}$ .

When  $(v_m, b_1)$  and  $(c_1, v_m)$  are two arcs of  $T_1$ , we see that if  $T_2 \neq F_4$  (possibly  $T_2 = F_3$ ; see Figure 8), then  $T \in \mathcal{P}$  with join vertices  $v_1, \dots, v_m, v_{m+1} := c_1, v_{m+2} := b_1$  and with new blocks  $A_m := A_m \setminus \{a_1, b_1, c_1\}$ ,  $B_m = \{c_1, v_m\}$ ,  $A_{m+1} = \{c_1\}$ ,  $B_{m+1} = \{b_1, c_1\}$ , and  $A_{m+2} = T_2 \setminus a_2$ ; if  $T_2 = F_4$  (see Figure 8), then  $T \in \mathcal{P}$  with join vertices  $v_1, \dots, v_m, v_{m+1} := c_1, v_{m+2} := b_1, v_{m+3} := z_3$  and with new blocks  $A_m := A_m \setminus \{a_1, b_1, c_1\}$ ,  $B_m = \{c_1, v_m\}$ ,  $A_{m+1} = \{c_1\}$ ,  $B_{m+1} = \{b_1, c_1\}$ ,  $A_{m+2} = \{b_1\}$ ,  $B_{m+2} = \{b_1, z_1, z_3, z_4\}$ , and  $A_{m+3} = \{z_2, z_3\}$ .

Combining the above observations, we conclude that  $T \in \mathcal{P}$ , thereby establishing the “only if” part.  $\blacksquare$

### 3 Minimax Relation

In this section we show that every Möbius-free strong tournament other than  $F_1$  and  $G_1$  satisfies the minimax relation on packing and covering feedback arc sets.

**Theorem 3.1.** *Let  $T = (V, A)$  be a Möbius-free strong tournament with  $|V| \geq 3$  and  $T \notin \{F_1, G_1\}$ . Then  $T$  is FAS Mengerian.*

As usual, we use  $\mathbb{Z}_+$  to denote the set of all nonnegative integers and use  $\mathbb{Z}_+^A$  to denote the set of vectors  $x = (x(a) : a \in A)$  whose coordinates belong to  $\mathbb{Z}_+$ . Let  $w \in \mathbb{Z}_+^A$ . A cycle  $C$  in  $T$  is called a *minimum cycle* of  $(T, w)$  if  $w(C) = \mu_w(T)$ . Let  $u$  and  $v$  be two vertices of  $T$ . A  $u - v$  path is a path from  $u$  to  $v$ . A  $u - v$  path is called *minimum* with respect to  $w$  (or simply  $w$ -*minimum*) if it has the minimum total weight among all  $u - v$  paths. An FAS packing of  $T$  with respect to  $w$  is also called a  $w$ -FAS packing.

By Theorem 2.7, every Möbius-free strong tournament  $T$  other than  $F_1$  and  $G_1$  can be depicted as in Figure 7. We shall prove Theorem 3.1 by induction on the number of vertical blocks in  $T$ ; the lemma below clearly yields the base statement.

**Lemma 3.2.** *Every tournament in  $\mathcal{Q}$  (see the paragraph succeeding Theorem 2.7) is FAS Mengerian.*

**Proof.** Let  $G = (V, A)$  be a tournament in  $\mathcal{Q}$ , let  $v$  be the center of  $G$ , and let  $H_1, H_2, \dots, H_k$  be the building blocks of  $G$  centered at  $v$ . We use  $\Omega$  to denote the set of all subscripts  $i$  with  $|H_i| = 4$  and use  $\Delta_i$  to denote the triangle  $H_i \setminus v$  for each  $i \in \Omega$ . Note that these triangles are pairwise vertex disjoint.

If  $v$  is a source or a sink of  $G$ , then the triangles  $\Delta_i$  are the only cycles in  $G$ . Thus  $G$  is trivially FAS Mengerian. So we assume hereafter that  $v$  is neither a source nor a sink of  $G$ .

Let  $w \in \mathbb{Z}_+^A$ . Our objective is to find a  $w$ -FAS packing in  $G$  of size  $r := \mu_w(G)$ . For this purpose, let  $X$  (resp.  $Y$ ) be the out-neighborhood (resp. in-neighborhood) of  $v$  in  $G$ , and let  $D$  be the digraph obtained from  $G$  by splitting  $v$  into a source  $s$  and a sink  $t$ , such that

- for each vertex  $x \in X$ , there is an arc  $sx$  in  $D$  with length  $w(sx) = w(vx)$ ;
- for each vertex  $y \in Y$ , there is an arc  $yt$  in  $D$  with length  $w(yt) = w(yv)$ ; and
- for each arc  $ab$  of  $G$  with  $v \notin \{a, b\}$ , there is an arc  $ab$  in  $D$  with length  $w(ab)$ .

Let  $\mathcal{C}$  be the collection of all cycles (directed) passing through  $v$  in  $G$ , and let  $r'$  be the minimum weight of a cycle in  $\mathcal{C}$ . Clearly,  $r' \geq r$ . We call a subset of arcs in  $G$  a  $\mathcal{C}$ -transversal if it intersects each cycle in  $\mathcal{C}$ . We also view  $\Delta_i$  for  $i \in \Omega$  as a triangle in  $D$  and view each arc of  $D$  as an arc of  $G$ .

From the construction of  $D$ , we see that

(1) there is a 1 – 1 correspondence between cycles in  $\mathcal{C}$  and  $s - t$  paths in  $D$ , and the shortest distance from  $s$  to  $t$  in  $D$  with respect to  $w$  is equal to  $r'$ .

For  $i = 1, 2, \dots, r$ , let  $U_i$  be the set of vertices at distance less than  $i$  from  $s$  in  $D$  with respect to  $w$ , and let  $C_i := \delta^+(U_i)$ . (Possibly there are arcs entering  $U_i$  in  $D$ , yet  $C_i$  consists of arcs leaving  $U_i$  only. So  $C_i$  is an  $s - t$  cut in  $D$ .) Observe that

(2) no  $C_i$  contains two or more arcs in  $\Delta_j$  for any  $j \in \Omega$  and

(3) each  $C_i$  corresponds to a  $\mathcal{C}$ -transversal in  $G$  by (1). Furthermore, each arc  $a$  of  $D$  is contained in at most  $w(a)$  of  $C_1, C_2, \dots, C_r$ .

Let us construct  $F_1, F_2, \dots, F_r$  from  $C_1, C_2, \dots, C_r$  by using the following algorithm.

Initially, set  $F_i := C_i$  for  $1 \leq i \leq r$ . While  $\Omega \neq \emptyset$ , do: take  $j \in \Omega$ , and add precisely one of the arcs  $e_{j,1}, e_{j,2}, e_{j,3}$  of  $\Delta_j$  to each  $F_i$  (if it contains no arc of  $\Delta_j$ ) to form a new  $F_i$  so that each  $e_{j,p}$  for  $1 \leq p \leq 3$  is contained in at most  $w(e_{j,p})$  of the resulting  $F_1, F_2, \dots, F_r$ . Set  $\Omega = \Omega - \{j\}$ .

Since  $\Delta_j$  is a triangle,  $w(e_{j,1}) + w(e_{j,2}) + w(e_{j,3}) \geq r$ . Thus the correctness of our algorithm is guaranteed by (2) and (3). From (3) and the structure of  $G$ , we further deduce that each  $F_i$  is an FAS of  $G$  and that each arc  $a$  of  $G$  is contained in at most  $w(a)$  members of  $\mathcal{F} := \{F_1, F_2, \dots, F_r\}$ . Therefore  $\mathcal{F}$  is a  $w$ -FAS packing of  $G$  having size  $r$ . ■

For convenience, we say that the  $w$ -FAS packing  $\mathcal{F}$  of size  $r$  output above is obtained by first performing breadth-first search for  $r$  steps in  $G$  from  $v$  and then eliminating triangles in  $G \setminus v$ , and say that  $F_i$  is the *depth- $i$  set* in  $\mathcal{F}$  from  $v$  for  $1 \leq i \leq r$ . Keep in mind that breadth-first

search employed in this paper always starts from a source (that is why we split  $v$  into a source and a sink as there are arcs entering and leaving it).

Let  $T = (V, A)$  be as described in Theorem 2.7, and let  $T^* = (V, E^*)$  be the subgraph of  $T$  arising from the vertical block  $A_1$  by adding all arcs  $ab$  of  $T$  with  $w(ab) > 0$ ,  $a \in V(A_1)$ , and  $b \notin V(A_1)$ . Note that  $T^* = T = A_1$  if  $m = 1$  and that  $T^*$  contains no arc in  $B_1$  except possibly  $v_1v_2$  when  $|B_1| = 4$ . For any collection  $\mathcal{F}$  of subsets of  $A$ , we use  $\mathcal{F} \cap E^*$  to denote the collection consisting of all nonempty  $F \cap E^*$  for  $F \in \mathcal{F}$ .

To prove Theorem 3.1, we shall actually show the existence of a  $w$ -FAS packing in  $T$  of size  $\mu_w(T)$  for all  $w \in \mathbb{Z}_+^A$  by induction on the number of vertical blocks. For this purpose, reducing arc weights while preserving the minimum total weight of a cycle whenever possible, it suffices to consider the weight functions  $w$  such that each arc  $e$  with  $w(e) > 0$  is contained in a minimum cycle of  $(T, w)$ . To make the induction work, what we establish is the following stronger statement.

**Theorem 3.3.** *Let  $T = (V, A)$  be a Möbius-free strong tournament with  $|V| \geq 3$  and  $T \notin \{F_1, G_1\}$ , and let  $w \in \mathbb{Z}_+^A$  such that each arc  $e$  with  $w(e) > 0$  is contained in a minimum cycle of  $(T, w)$ . Then  $T$  has a  $w$ -FAS packing  $\mathcal{F}$  of size  $\mu_w(T)$ , such that  $\mathcal{F} \cap E^*$  can be obtained by first performing breadth-first search for  $|\mathcal{F} \cap E^*|$  steps in  $T^*$  from  $v_1$  and then eliminating triangles in  $A_1 \setminus v_1$ .*

**Proof.** By Theorem 2.7,  $T$  can be depicted as in Figure 7, with vertical blocks  $A_1, A_2, \dots, A_m$ , horizontal blocks  $B_1, B_2, \dots, B_{m-1}$ , and join vertices  $v_1, v_2, \dots, v_m$ . We apply induction on  $m$ . Since each  $A_i \in \mathcal{Q}$ , the induction base  $m = 1$  follows instantly from Lemma 3.2. So we proceed to the induction step and assume that  $m \geq 2$  and that the statement holds for  $m - 1$ .

Let us first make some simple observations about the weight function  $w$ .

(1) For any arc  $uv$  and any path  $P$  from  $u$  to  $v$  in  $T$ , we have  $w(uv) \leq w(P)$ .

Assume that contrary:  $w(uv) > w(P)$ . By hypothesis,  $uv$  is contained in a minimum cycle  $C$  of  $(T, w)$ . Let  $D$  be the multiset union of  $P$  and  $C[v, u]$  (that is, if an arc is contained in both  $P$  and  $C[v, u]$ , then it appears twice in  $D$ ). Clearly,  $D$  is an Eulerian digraph with  $w(D) < w(C)$ . Let  $C'$  be a directed cycle contained in  $D$ . Then  $w(C') \leq w(D) < w(C)$ , contradicting the minimality assumption on  $C$ .

From (1) it is clear that

(2) for any minimum cycle  $C$  of  $(T, w)$  and any chord  $uv$  of  $C$ , the cycle arising from  $C$  by replacing  $C[u, v]$  with  $uv$  is also minimum. So  $w(uv) = w(C[u, v])$ .

(3) There is a path in  $A_1$  from  $v_1$  to each vertex in  $A_1 \setminus v_1$ . So  $v_1$  is not a sink of  $A_1$ .

To justify this, note from the structural description (see Theorem 2.7) that  $A_1 \setminus v_1$  contains either a source  $u$  or a triangle  $\Delta$  such that no arc enters it (in  $A_1 \setminus v_1$ ). In the former case,  $v_1u$  is an arc of  $T$ ; in the latter case, there is at least one arc from  $v_1$  to  $\Delta$  in  $T$ , for otherwise  $\delta(u)$  or  $\delta(\Delta)$  would be a dicut in  $T$ , contradicting the strong connectivity of  $T$ . Thus (3) holds.

(4) Let  $ab$  be an arc in  $T$  with  $w(ab) > 0$ ,  $a \in V(A_1 \setminus v_1)$ , and  $b \notin V(A_1 \cup B_1 \setminus v_2)$ , and let  $P$  be a minimum  $v_1 - a$  path in  $A_1$  (see (3)). If  $v_1b$  is an arc of  $T$ , then  $w(P) + w(ab) = w(v_1b)$ . (Possibly  $b = v_2$  when  $|B_1| = 4$ .)

To justify this, let  $C$  be a minimum cycle of  $(T, w)$  containing  $ab$ . From the structural description of  $T$ , we see that  $C$  passes through  $v_1$  and that  $C[v_1, a]$  is fully contained in  $A_1$ . By the minimality assumptions on  $P$  and  $C$ , we obtain  $w(P) = w(C[v_1, a])$ . In view of (2),

$w(C[v_1, b]) = w(v_1b)$ . Hence  $w(P) + w(ab) = w(C[v_1, a]) + w(ab) = w(C[v_1, b]) = w(v_1b)$ , as desired.

Let  $T_1 = (V_1, E_1)$  be the subtournament of  $T$  induced by all vertices in  $A_1 \cup B_1$ . Then  $T_1 \in \mathcal{Q}$ . Let  $T_2 = (V_2, E_2)$  be the subtournament of  $T$  induced by all vertices outside  $A_1 \setminus v_1$ , and let  $A'_2$  be the subtournament of  $T$  induced by all vertices in  $A_2 \cup B_1$ . By Theorem 2.7, we have

(5)  $T_i$  is a Möbius-free strong tournament with  $|V_i| \geq 3$  and  $T_i \notin \{F_1, G_1\}$  for  $i = 1, 2$ . Furthermore,  $A'_2, A_3, \dots, A_m$  are the vertical blocks of  $T_2$ .

In the remainder of our proof, we reserve  $u_1$  for the vertex in  $B_1 \setminus \{v_1, v_2\}$  if  $|B_1| = 3$ , and reserve  $u_1$  and  $u_2$  for the two vertices in  $B_1 \setminus \{v_1, v_2\}$  if  $|B_1| = 4$ , with  $u_1u_2 \in A$ . Moreover, we reserve  $R_1$  for a minimum  $v_2 - v_1$  path in  $B_1$  with respect to  $w$ , having the fewest arcs. By (1), we obtain  $R_1 = v_2v_1$  if  $|B_1| \leq 3$  and  $R_1 = v_2u_1v_1$  or  $v_2u_2v_1$  if  $|B_1| = 4$ . Write  $r := \mu_w(T)$ . The statement below follows instantly from (2).

(6) Each arc  $e$  in  $B_1$  with  $w(e) > 0$  is contained in a cycle of  $T_i$  with weight  $r$  for  $i = 1$  or  $2$  (but not necessarily both). Each arc  $e$  in  $T_i$  but outside  $B_1$  with  $w(e) > 0$  is contained in a cycle of  $T_i$  with weight  $r$  for  $i = 1, 2$ . Furthermore, if  $|B_1| = 4$ , then the arc  $u_iv_1$  with  $w(u_iv_1) > 0$  is contained in a cycle of  $T_1$  with weight  $r$ , and the arc  $v_2u_i$  with  $w(v_2u_i) > 0$  is contained in a cycle of  $T_2$  with weight  $r$  for  $i = 1, 2$ .

(7) For each vertex  $a$  in  $A_1 \setminus v_1$  with  $w(av_2) > 0$ , the path  $av_2R_1$  is contained in a cycle of  $T_1$  with weight  $r$ . For each vertex  $b$  outside  $A_1 \cup B_1 \setminus v_2$  with  $w(v_1b) > 0$ , the path  $R_1v_1b$  is contained in a cycle of  $T_2$  with weight  $r$ .

We only establish the second half here, as the proof of the first half goes along the same line. By (6), arc  $v_1b$  is contained in a cycle  $C$  of  $T_2$  with weight  $r$ . Since  $\delta(B_1 \setminus v_2)$  forms a dicut in  $T_2 \setminus v_2$ , cycle  $C$  must pass through  $v_2$ . It follows that  $C[v_2, v_1]$  is fully contained in  $B_1$ . Let  $C'$  be obtained from  $C$  by replacing  $C[v_2, v_1]$  with  $R_1$ . Then  $C'$  is a cycle of  $T_2$  with weight  $r$  and contains the path  $R_1v_1b$ . So (7) is justified.

(8) If  $R_1$  is not contained in any cycle of  $T_1$  with weight  $r$ , then  $w(ab) = 0$  for any  $a \in V(A_1)$  if  $|B_1| = 4$  and  $a \in V(A_1 \setminus v_1)$  if  $|B_1| \leq 3$  and  $b \notin V(A_1 \cup B_1 \setminus v_2)$ . If  $R_1$  is not contained in any cycle of  $T_2$  with weight  $r$ , then  $w(ab) = 0$  for any  $a \in V(A_1)$  and  $b \notin V(A_1 \cup B_1 \setminus v_2)$  if  $|B_1| = 4$  and  $b \notin V(A_1 \cup B_1)$  if  $|B_1| \leq 3$ .

Suppose on the contrary that  $w(ab) > 0$  for some  $a \in V(A_1)$  if  $|B_1| = 4$  or  $a \in V(A_1 \setminus v_1)$  if  $|B_1| = 3$  and  $b \notin V(A_1 \cup B_1 \setminus v_2)$ . Let  $C$  be a minimum cycle of  $(T, w)$  containing  $ab$ . From Theorem 2.7 we see that  $C$  passes through  $v_1$  and  $v_2$  and that  $C[v_2, v_1]$  is fully contained in  $B_1$ . If  $a \in V(A_1 \setminus v_1)$ , then  $av_2$  is  $ab$  or a chord of  $C$ . By (2), we have  $w(av_2) = w(C[a, v_2]) \geq w(ab) > 0$ . It follows from (7) that  $R_1$  is contained in a cycle of  $T_1$  with weight  $r$ , a contradiction. So we assume that  $a = v_1$  and  $|B_1| = 4$ . By (2), the cycle arising from  $C[v_2, v_1]$  by adding  $v_1v_2$  is a minimum cycle of  $(T, w)$ . Therefore  $v_2R_1v_1v_2$  is also a cycle of  $T_1$  with weight  $r$ , a contradiction again. The second half of the statement can be established similarly.

For  $i = 1, 2$ , define  $w_i \in \mathbb{Z}_+^{E_i}$  to be the weight function obtained from  $w|_{E_i}$  by reducing the weights of arcs in  $B_1$ , if necessary, so that  $\mu_{w_i}(T_i) = r$  and that each arc  $e$  in  $T_i$  with  $w_i(e) > 0$  is contained in a minimum cycle of  $(T_i, w_i)$  (see (6)). We point out that  $w_1(v_1v_2) = w_2(v_1v_2) = w(v_1v_2)$  when  $|B_1| = 4$ ; we postpone giving its proof till this case is discussed (see (23)), as this observation has nothing to do with the case when  $|B_1| \leq 3$ .

Let  $T_1^* = (V, E_1^*)$  be the subgraph of  $T$  obtained from  $T_1 = (V_1, E_1)$  by adding all arcs  $ab$  with  $w(ab) > 0$ ,  $a \in V(A_1)$ , and  $b \notin V(A_1 \cup B_1)$ , and define  $w_1(ab) = w(ab)$  for each such arc  $ab$ . Let  $T_2^* = (V_2, E_2^*)$  be the subgraph of  $T_2 = (V_2, E_2)$  arising from block  $A_2'$  by adding all arcs  $ab$  with  $w(ab) > 0$ ,  $a \in V(A_2')$ , and  $b \notin V(A_2')$ . For ease of description, we color each arc  $v_1b$ , with  $w(v_1b) > 0$  and  $b \notin V(A_1 \cup B_1 \setminus v_2)$ , by blue. (Possibly  $b = v_2$  when  $|B_1| = 4$ ). From (4) and the proof of Lemma 3.2, we see that

(9)  $T_1^*$  has a  $w_1$ -FAS packing  $\mathcal{F}_1$  of size  $r$ , obtained by first performing breadth-first search (with respect to the weight function  $w_1$ ) for  $r$  steps from  $v_1$  in  $T_1^*$  and then eliminating triangles in  $A_1 \setminus v_1$ , such that each blue arc  $e$  is contained in precisely  $w(e)$  members of  $\mathcal{F}_1$ .

Using (5) and induction hypothesis, we deduce that

(10)  $T_2$  has a  $w_2$ -FAS packing  $\mathcal{F}_2$  of size  $r$ , such that  $\mathcal{F}_2 \cap E_2^*$  can be obtained by first performing breadth-first search (with respect to the weight function  $w_2$ ) for  $|\mathcal{F}_2 \cap E_2^*|$  steps in  $T_2^*$  from  $v_2$  and then eliminating triangles in  $A_2' \setminus v_2$ .

We shall produce a  $w$ -FAS packing  $\mathcal{F}$  of  $T$  having size  $r$  by gluing members of  $\mathcal{F}_1$  together with those of  $\mathcal{F}_2$ , possibly with slight modification. For  $i = 1, 2$ , let  $\mathcal{F}_i = \{F_{i,1}, F_{i,2}, \dots, F_{i,r}\}$ , where  $F_{i,j}$  is the depth- $j$  set in  $\mathcal{F}_1$  from  $v_1$ , and  $F_{i,j} \cap E_2^*$  is the depth- $j$  set in  $\mathcal{F}_2 \cap E_2^*$  from  $v_2$ . We color each  $F_{i,j}$  containing a blue arc also by blue. Observe that no arc, except blue ones and those in  $B_1$ , is shared by members of  $\mathcal{F}_1$  and members of  $\mathcal{F}_2$ . So, naturally, in our proof blue members of  $\mathcal{F}_1$  will be glued together with blue members of  $\mathcal{F}_2$ . Once the members of  $\mathcal{F}$  containing blue arcs are determined, the members containing arcs  $ab$  with  $a \in V(A_1 \setminus v_1)$  and  $b \notin V(A_1 \cup B_1 \setminus v_2)$  will be determined accordingly by (4).

Depending on the size of  $B_1$ , we distinguish between two cases.

**Case 1.**  $|B_1| \leq 3$ .

We may assume that  $|B_1| = 3$ , because this situation properly contains the one when  $|B_1| = 2$ . Let  $q := w(v_2v_1)$ ,  $s := w(u_1v_1)$ , and  $t := w(v_2u_1)$ . In view of (1), we have  $q \leq s + t$ .

(11) If  $s > 0$  and  $u_1v_1$  is not contained in a cycle of  $T_1$  having weight  $r$  with respect to the weight function  $w$ , then  $q = s + t$ . Furthermore,  $v_2v_1$  is not contained in a cycle of  $T_1$  having weight  $r$  with respect to  $w$  either.

By (6),  $u_1v_1$  is contained in a cycle  $C$  of  $T_2$  having weight  $r$  with respect to  $w$ . Clearly,  $C$  passes through  $v_2u_1$ . It follows instantly from (2) that  $q = s + t$ . Assume on the contrary that  $v_2v_1$  is contained in a cycle  $Q$  of  $T_1$  having weight  $r$  with respect to  $w$ . Let  $Q'$  be the cycle obtained from  $Q$  by replacing  $v_2v_1$  with the path  $v_2u_1v_1$ . Then  $Q'$  has weight  $r$  and contains  $u_1v_1$ , a contradiction. So (11) is justified.

Similarly, the following statement holds.

(12) If  $t > 0$  and  $v_2u_1$  is not contained in a cycle of  $T_2$  having weight  $r$  with respect to the weight function  $w$ , then  $q = s + t$ . Furthermore,  $v_2v_1$  is not contained in a cycle of  $T_2$  having weight  $r$  with respect to  $w$  either.

Let  $E_1'$  be the arc set obtained from  $E_1^*$  by deleting arcs in  $B_1$ , let  $E_2'$  be the arc set obtained from  $E_2$  by deleting arcs in  $B_1$ , and let  $K_{i,j}$  be the restriction of  $F_{i,j}$  to  $E_i'$  for  $i = 1, 2$  and  $1 \leq j \leq r$ .

(13) Let us modify  $K_{i,j}$ 's as follows:

- add arc  $v_2v_1$  to  $K_{1,j}$  for  $r - q + 1 \leq j \leq r$ ;
- add arc  $u_1v_1$  to  $K_{1,j}$  for  $r - s + 1 \leq j \leq r$ ;

- add arc  $v_2u_1$  to  $K_{1,j}$  for  $r - q + 1 \leq j \leq r - s$ ;
- add arc  $v_2v_1$  to  $K_{2,j}$  for  $1 \leq j \leq q$ ;
- add arc  $v_2u_1$  to  $K_{2,j}$  for  $1 \leq j \leq t$ ; and
- add arc  $u_1v_1$  to  $K_{2,j}$  for  $t + 1 \leq j \leq q$ .

We use  $F'_{i,j}$  to denote the resulting  $K_{i,j}$ .

(14)  $\mathcal{F}'_1 := \{F'_{1,1}, F'_{1,2}, \dots, F'_{1,r}\}$  is a  $w$ -FAS packing of  $T_1^*$ , and  $\mathcal{F}'_2 := \{F'_{2,1}, F'_{2,2}, \dots, F'_{2,r}\}$  is a  $w$ -FAS packing of  $T_2$ .

To justify this, recall that each arc  $e \in F_{1,j}$  satisfies  $w_1(e) > 0$  and that each arc  $e$  of  $T_1$  with  $w_1(e) > 0$  is contained in a cycle of  $T_1$  with weight  $r$ . From (9) and breadth-first search we deduce that

- $F_{1,j}$  contains  $v_2v_1$  iff  $r - w_1(v_2v_1) + 1 \leq j \leq r$ ;
- $F_{1,j}$  contains  $u_1v_1$  iff  $r - w_1(u_1v_1) + 1 \leq j \leq r$ ; and
- $F_{1,j}$  contains  $v_2u_1$  iff  $r - w_1(v_2u_1) + 1 \leq j \leq r - w_1(u_1v_1)$ .

Since  $q \geq w_1(v_2v_1)$ ,  $s \geq w_1(u_1v_1)$ , and  $t \geq w_1(v_2u_1)$ , we deduce that if  $F_{1,j}$  contains  $v_2v_1$ , then so does  $F'_{1,j}$ , and if  $F_{1,j}$  contains  $u_1v_1$ , then so does  $F'_{1,j}$ . Moreover, if  $F_{1,j}$  contains  $v_2u_1$ , then  $F'_{1,j}$  contains  $v_2u_1$  or  $u_1v_1$ . Note that each cycle of  $T_1$  containing  $v_2u_1$  must pass through  $u_1v_1$ . Since each  $F_{1,j}$  is an FAS of  $T_1^*$ , so is  $F'_{1,j}$ . From (13) it is clear that  $\mathcal{F}'_1$  is a  $w$ -FAS packing of  $T_1^*$ . Similarly, we can prove that  $\mathcal{F}'_2$  is a  $w$ -FAS packing of  $T_2$ .

(15) If  $F'_{2,j} \neq F_{2,j}$  for some  $j$  with  $1 \leq j \leq r$ , then  $w(ab) = 0$  for any  $a \in V(A_1)$  and  $b \notin V(A_1 \cup B_1)$ . In particular, there is no blue arc in  $T$ .

To justify this, note from (1), (10) and breadth-first search that  $F_{2,j}$  contains  $v_2v_1$  iff  $1 \leq j \leq w_2(v_2v_1)$ ,  $F_{2,j}$  contains  $v_2u_1$  iff  $1 \leq j \leq w_2(v_2u_1)$ , and  $F_{2,j}$  contains  $u_1v_1$  iff  $w_2(v_2u_1) + 1 \leq j \leq w_2(v_2v_1)$ . Since  $F'_{2,j} \neq F_{2,j}$  for some  $j$  with  $1 \leq j \leq r$ , we deduce from (13) that  $w_2(v_2v_1) < q$  or  $w_2(v_2u_1) < t$ . From (12) we further conclude that the inequality  $w_2(v_2v_1) < q$  must hold. Thus (15) follows instantly from (8).

From (9), (10), (13) and (15) we see that

(16)  $F'_{1,j}$  contains a blue arc iff  $F'_{2,j+q}$  contains it.

Define

$$(17) F_j := \begin{cases} F'_{1,j} \cup F'_{2,j+q} & \text{if } 1 \leq j \leq r - q; \\ F'_{1,j} \cup F'_{2,j+q-r} & \text{if } r - q + 1 \leq j \leq r. \end{cases}$$

(18) For  $F_j$ 's defined in (17), the following statements hold:

- $F_j$  contains  $v_2v_1$  iff  $r - q + 1 \leq j \leq r$ ;
- $F_j$  contains  $u_1v_1$  iff  $r - s + 1 \leq j \leq r$ ; and
- $F_j$  contains  $v_2u_1$  iff  $r - q + 1 \leq j \leq \min\{r, r - q + t\}$  or  $1 \leq j \leq \max\{0, t - q\}$ .

To justify this, note from (17) that  $F'_{1,j}$  is a subset of  $F_j$  for  $1 \leq j \leq r$  and from (13) that

(18.1)  $F'_{1,j}$  contains  $v_2v_1$  iff  $r - q + 1 \leq j \leq r$ , and  $F'_{2,k}$  contains  $v_2v_1$  iff  $1 \leq k \leq q$ ;

(18.2)  $F'_{1,j}$  contains  $u_1v_1$  iff  $r - s + 1 \leq j \leq r$ , and  $F'_{2,k}$  contains  $u_1v_1$  iff  $t + 1 \leq k \leq q$ ;

(18.3)  $F'_{1,j}$  contains  $v_2u_1$  iff  $r - q + 1 \leq j \leq r - s$ , and  $F'_{2,k}$  contains  $v_2u_1$  iff  $1 \leq k \leq t$ .

First, let  $k$  be a subscript with  $v_2v_1 \in F'_{2,k}$ . Then  $1 \leq k \leq q$  by (18.1). Let  $j$  be the subscript with  $k = j + q - r$ . Then  $j = r - q + k$ . Thus  $r - q + 1 \leq j \leq r$  and hence  $F'_{2,k}$  is a subset of  $F_j$  by (17). Combining this with (18.1) (as  $F'_{1,j} \subseteq F_j$ ), we see that  $F_j$  contains  $v_2v_1$  iff  $r - q + 1 \leq j \leq r$ .

Second, let  $k$  be a subscript with  $u_1v_1 \in F'_{2,k}$ . Then  $t + 1 \leq k \leq q$  by (18.2). Let  $j$  be the subscript with  $k = j + q - r$ . Then  $j = r - q + k$ . Thus  $r - q + t + 1 \leq j \leq r$ . It follows from

(17) that  $F'_{2,k}$  is a subset of  $F_j$ . By (1),  $s + t \geq q$ . So  $r - q + t + 1 \geq r - s + 1$  and hence  $r - s + 1 \leq j \leq r$ . Combining this with (18.2) (as  $F'_{1,j} \subseteq F_j$ ), we see that  $F_j$  contains  $u_1v_1$  iff  $r - s + 1 \leq j \leq r$ .

Finally, let  $k$  be a subscript with  $v_2u_1 \in F'_{2,k}$ . Then  $1 \leq k \leq t$  by (18.3). When  $1 \leq k \leq \min\{q, t\}$ , let  $j$  be the subscript with  $k = j + q - r$ . Then  $j = r - q + k$ . So  $r - q + 1 \leq j \leq \min\{r, r - q + t\}$  and thus  $F'_{2,k}$  is a subset of  $F_j$  by (17). When  $\min\{q, t\} + 1 \leq k \leq t$  (equivalently  $q + 1 \leq k < t$ ), let  $j$  be the subscript with  $k = j + q$ . Then  $j = k - q$ . Thus  $1 \leq j \leq t - q$  and hence  $F'_{2,k}$  is a subset of  $F_j$  by (17). Therefore, there exists a subscript  $k$  with  $v_2u_1 \in F'_{2,k} \subseteq F_j$  iff  $r - q + 1 \leq j \leq \min\{r, r - q + t\}$  or  $1 \leq j \leq \max\{0, t - q\}$ . Combining this with (18.3) (as  $F'_{1,j} \subseteq F_j$ ), we see that  $F_j$  contains  $v_2u_1$  iff  $r - q + 1 \leq j \leq \min\{r, r - q + t\}$  or  $1 \leq j \leq \max\{0, t - q\}$ , because  $r - s \leq \min\{r, r - q + t\}$  (recall that  $s + t \geq q$  by (1)). This establishes (18).

In view of (16)-(18), we obtain

(19) each arc  $e$  of  $T$  is contained in at most  $w(e)$  members of  $\mathcal{F} := \{F_1, F_2, \dots, F_r\}$ .

Let us show that

(20) each  $F_j$ , with  $1 \leq j \leq r$ , is an FAS of  $T$ .

For this purpose, let  $C$  be an arbitrary cycle in  $T$ . Clearly,  $F_j$  intersects  $C$  if  $C$  is a cycle of  $T_1$  or a cycle of  $T_2$  by (14). So we assume that  $C$  is not fully contained in  $T_i$  for  $i = 1, 2$ .

Consider the subcase when  $u_1$  is outside  $C$ . Now  $C$  contains an arc  $ab$  with  $a \in V(A_1 \setminus v_1)$  and  $b \notin V(A_1 \cup B_1)$ . From Theorem 2.7 we deduce that  $C$  passes through  $v_1$  and  $C[v_1, a]$  is fully contained in  $A_1$ . Let  $C'$  be the cycle arising from  $C$  by replacing  $C[v_1, b]$  with  $v_1b$ , and let  $F'_{2,k}$  be the member of  $\mathcal{F}_2$  contained in  $F_j$ . Then  $C'$  is fully contained in  $T_2$  and intersects  $F'_{2,k}$ . If  $F'_{2,k}$  intersects  $C'[b, v_1] = C[b, v_1]$ , then  $F_j$  intersects  $C$ . So we assume that  $F'_{2,k}$  contains  $v_1b$  and hence  $w(v_1b) > 0$ , indicating that  $v_1b$  is a blue arc. By (8),  $w_2(v_2v_1) = w(v_2v_1) = q$ . Thus, by (16) and (17),  $k = j + q$  and  $F'_{1,j}$  contains  $v_1b$  as well. In view of (1),  $w_1(C[v_1, b]) = w(C[v_1, b]) \geq w(v_1b)$ . From the construction of  $\mathcal{F}_1$  using depth-first search, we see that  $F'_{1,j}$  intersects  $C[v_1, b]$ . So  $F_j$  intersects  $C$ .

It remains to consider the subcase when  $C$  contains  $u_1$ . Assume first that  $v_2u_1v_1$  is a segment of  $C$ . Let  $C'$  be obtained from  $C$  by replacing  $v_2u_1v_1$  with  $v_2v_1$ . As observed in the preceding paragraph,  $F_j$  intersects  $C'$ . If  $v_2v_1 \notin F_j$ , then  $F_j$  intersects  $C'[v_1, v_2]$  and hence  $C$ ; otherwise,  $v_2v_1 \in F_j$ , so  $r - q + 1 \leq j \leq r$  by (18). Since  $s + t \geq q$  by (1), we have  $r - q + t \geq r - s$ . Hence  $r - q + 1 \leq j \leq \min\{r, r - q + t\}$  or  $r - s + 1 \leq j \leq r$ . It follows from (18) that  $F_j$  contains  $v_2u_1$  or  $u_1v_1$ . Therefore  $F_j$  intersects  $C$ .

Next, we assume that  $C$  has a segment  $au_1b$ , where  $a \in V(A_1 \setminus v_1)$  and  $b \notin V(A_1 \cup B_1)$ . Note that  $v_2v_1$  is contained in  $C$  and  $C[v_1, a]$  is fully contained in  $A_1$ . Let  $C'$  be obtained from  $C$  by replacing  $C[v_2, u_1]$  with  $v_2u_1$ . Then  $C'$  is fully contained in  $T_2$ . So  $F_j$  intersects  $C'$ . If  $v_2u_1 \notin F_j$ , then  $F_j$  intersects  $C'[u_1, v_2]$  and hence  $C$ ; otherwise,  $v_2u_1 \in F_j$ , so  $r - q + 1 \leq j \leq \min\{r, r - q + t\}$  or  $1 \leq j \leq \max\{0, t - q\}$  by (18). If  $t \leq q$ , then  $r - q + 1 \leq j \leq r - q + t \leq r$ . Thus  $F_j$  contains  $v_2v_1$  by (18) and hence intersects  $C$ . Suppose  $t > q$ . Since  $(r - s) + q \geq t$  by (11) or (1) (when  $s > 0$  and  $u_1v_1$  is contained in a cycle  $Q$  of  $T_1$  having weight  $r$ , consider the path  $v_2v_1Q[v_1, u_1]$ , which has weight  $(r - s) + q$ ), we obtain  $r - s \geq t - q$ . Hence  $r - q + 1 \leq j \leq r$  or  $1 \leq j \leq r - s$ . It follows from (18) that either  $F_j$  contains  $v_2v_1$  or  $F'_{1,j}$  (and hence  $F_j$ ) intersects  $C[v_1, u_1]$  by (9). This establishes (20).

Combining (19) with (20), we conclude that  $\mathcal{F}$  is a  $w$ -FAS packing of  $T$  having size  $r$ . From

(9) and (15)-(17), it is clear that  $\mathcal{F} \cap E^*$  is obtained by first performing breadth-first search for  $|\mathcal{F} \cap E^*|$  steps in  $T^*$  from  $v_1$  and then eliminating triangles in  $A_1 \setminus v_1$ .

**Case 2.**  $|B_1| = 4$ .

Observe that

(21) arc  $v_1v_2$  is contained in only three cycles,  $v_1v_2u_1v_1$ ,  $v_1v_2u_2v_1$ , and  $v_1v_2u_1u_2v_1$ , of  $T$ , and  $w(v_1v_2u_iv_1) \leq w(v_1v_2u_1u_2v_1)$  for  $i = 1, 2$  by (1).

(22)  $w(av_2) \leq w(v_1v_2)$  for any vertex  $a$  in  $A_1 \setminus v_1$ , and  $w(v_1b) \leq w(v_1v_2)$  for any vertex  $b$  outside  $A_1 \cup B_1 \setminus v_2$ .

We only prove the first half of this statement, as the proof of the second half does along the same line. If  $w(av_2) = 0$ , then trivially  $w(av_2) \leq w(v_1v_2)$ . So we assume that  $w(av_2) > 0$ . By (7), the path  $av_2R_1$  is contained in a cycle  $C$  of  $T_1$  having weight  $r$  with respect to the weight function  $w$ . By (1), we have  $w(C[v_1, v_2]) = w(v_1v_2)$ . It follows that  $w(av_2) \leq w(v_1v_2)$ . This establishes (22).

Let  $p := w(u_1u_2)$ ,  $q := w(v_1v_2)$ ,  $s_i := w(u_iv_1)$ , and  $t_i := w(v_2u_i)$  for  $i = 1, 2$ .

(23)  $q = w_1(v_1v_2) = w_2(v_1v_2)$ ,  $s_i = w_1(u_iv_1)$ , and  $t_i = w_2(v_2u_i)$  for  $i = 1, 2$ . Furthermore, if  $p > 0$ , then either  $p + s_2 = s_1$  or  $t_1 + p = t_2$ . If  $q > 0$ , then  $v_1v_2R_1v_1$  is a cycle having weight  $r$  with respect to the weight function  $w$ .

From (21) and (6) it follows immediately that  $q = w_1(v_1v_2) = w_2(v_1v_2)$ ,  $s_i = w_1(u_iv_1)$ , and  $t_i = w_2(v_2u_i)$  for  $i = 1, 2$ . To show the statements concerning  $p$ , let  $C$  be a cycle in  $T_i$  containing  $u_1u_2$  and having weight  $r$  with respect to the weight function  $w$  for  $i = 1$  or  $2$ ; such  $C$  exists by (6). Since  $u_2v_1$  is the only arc leaving  $u_2$  in  $T_1$ , and  $v_2u_1$  is the only arc entering  $u_1$  in  $T_2$ , cycle  $C$  contains  $u_2v_1$  or  $v_2u_1$ . Thus  $p + s_2 = s_1$  or  $t_1 + p = t_2$  by (2). If  $q > 0$ , then  $v_1v_2$  is contained in a cycle having weight  $r$  with respect to the weight function  $w$ . From (21) we deduce that  $v_1v_2R_1v_1$  has weight  $r$  with respect to  $w$ . So (23) is established.

We proceed by considering two subcases.

**Subcase 2.1.**  $s_i + t_i + q = r$  for  $i = 1$  or  $2$ .

From (4) and (7)-(10) we see that

(24)  $F_{1,j}$  contains a blue arc  $v_1b$  iff so does  $F_{2,j+r-q}$ . (Hence  $F_{1,j}$  is colored blue iff so is  $F_{2,j+r-q}$ .) Furthermore,  $F_{1,j}$  contains a blue arc iff  $1 \leq j \leq q$  by (22) and (23).

Define

$$(25) F_j := \begin{cases} F_{1,j} \cup F_{2,j+r-q} & \text{if } 1 \leq j \leq q; \\ F_{1,j} \cup F_{2,j-q} & \text{if } q+1 \leq j \leq r. \end{cases}$$

Thus each blue set in  $\mathcal{F}_1$  is glued together with the corresponding blue set in  $\mathcal{F}_2$  (see (24)), if any.

(26) For  $F_j$ 's defined in (25), the following statements hold:

- $F_j$  contains  $v_1v_2$  iff  $1 \leq j \leq q$ ;
- $F_j$  contains  $u_iv_1$  iff  $r - s_i + 1 \leq j \leq r$  for  $i = 1, 2$ ;
- $F_j$  contains  $v_2u_i$  iff  $q + 1 \leq j \leq \min\{r, q + t_i\}$  or  $1 \leq j \leq \max\{0, t_i - r + q\}$  for  $i = 1, 2$ ; and
- $F_j$  contains  $u_1u_2$  iff  $t_1 + q + 1 \leq j \leq \min\{r, t_2 + q\}$  or  $1 \leq j \leq \max\{0, t_2 - r + q\}$  when  $s_1 + t_1 + q = r$  and iff  $r - s_1 + 1 \leq j \leq r - s_2$  when  $s_2 + t_2 + q = r$ .

To justify this, note from (6), (9), (10) and (23) that

(26.1)  $F_{1,j}$  contains  $v_1v_2$  iff  $1 \leq j \leq q$ , and  $F_{2,k}$  contains  $v_1v_2$  iff  $r - q + 1 \leq k \leq r$ ;

(26.2)  $F_{1,j}$  contains  $u_i v_1$  iff  $r - s_i + 1 \leq j \leq r$ , and  $F_{2,k}$  contains  $u_i v_1$  iff  $t_i + 1 \leq k \leq r - q$  for  $i = 1, 2$ ;

(26.3)  $F_{1,j}$  contains  $v_2 u_i$  iff  $q + 1 \leq j \leq r - s_i$ , and  $F_{2,k}$  contains  $v_2 u_i$  iff  $1 \leq k \leq t_i$  for  $i = 1, 2$ ;

(26.4)  $F_{1,j}$  contains  $u_1 u_2$  iff  $r - s_1 + 1 \leq j \leq r - s_2$ , and  $F_{2,k}$  contains  $u_1 u_2$  iff  $t_1 + 1 \leq k \leq t_2$ .

First, let  $k$  be a subscript with  $v_1 v_2 \in F_{2,k}$ . Then  $r - q + 1 \leq k \leq r$  by (26.1). Let  $j$  be the subscript with  $k = j + r - q$ . Then  $j = k - r + q$ . Thus  $1 \leq j \leq q$  and hence  $F_{2,k}$  is a subset of  $F_j$  by (25). Combining this with (26.1) (as  $F_{1,j} \subseteq F_j$ ), we see that  $F_j$  contains  $v_1 v_2$  iff  $1 \leq j \leq q$ .

Second, let  $k$  be a subscript with  $u_i v_1 \in F_{2,k}$ . Then  $t_i + 1 \leq k \leq r - q$  by (26.2). Let  $j$  be the subscript with  $k = j - q$ . Then  $j = k + q$ . Thus  $t_i + q + 1 \leq j \leq r$ . Since  $s_i + t_i + q \geq r$ , we have  $t_i + q + 1 \geq r - s_i + 1$  and hence  $r - s_i + 1 \leq j \leq r$ . Combining this with (26.2) (as  $F_{1,j} \subseteq F_j$ ), we see that  $F_j$  contains  $u_i v_1$  iff  $r - s_i + 1 \leq j \leq r$ .

Third, let  $k$  be a subscript with  $v_2 u_i \in F_{2,k}$ . Then  $1 \leq k \leq t_i$  by (26.3). When  $1 \leq k \leq \min\{r - q, t_i\}$ , let  $j$  be the subscript with  $k = j - q$ . Then  $j = q + k$ . So  $q + 1 \leq j \leq \min\{r, q + t_i\}$ . Hence  $F_{2,k}$  is a subset of  $F_j$  by (25). When  $\min\{r - q, t_i\} + 1 \leq k \leq t_i$  (equivalently  $r - q + 1 \leq k \leq t_i$ ), let  $j$  be the subscript with  $k = j + r - q$ . Then  $j = k - r + q$ . Thus  $1 \leq j \leq t_i - r + q \leq q$  and hence  $F_{2,k}$  is a subset of  $F_j$  by (25). Therefore, there exists a subscript  $k$  with  $v_2 u_i \in F_{2,k} \subseteq F_j$  iff  $q + 1 \leq j \leq \min\{r, q + t_i\}$  or  $1 \leq j \leq \max\{0, t_i - r + q\}$ . Combining this with (26.3) (as  $F_{1,j} \subseteq F_j$ ), we see that  $F_j$  contains  $v_2 u_i$  iff  $q + 1 \leq j \leq \min\{r, q + t_i\}$  or  $1 \leq j \leq \max\{0, t_i - r + q\}$ , because  $s_i + t_i + q \geq r$ , which implies  $r - s_i \leq q + t_i$ .

Finally, let  $k$  be a subscript with  $u_1 u_2 \in F_{2,k}$ . Then  $t_1 + 1 \leq k \leq t_2$  by (26.4). When  $t_1 + 1 \leq k \leq \min\{r - q, t_2\}$ , let  $j$  be the subscript with  $k = j - q$ . Then  $j = k + q$ . Thus  $t_1 + q + 1 \leq j \leq \min\{r, t_2 + q\}$ . Hence  $F_{2,k}$  is a subset of  $F_j$  by (25). When  $\min\{r - q, t_2\} + 1 \leq k \leq t_2$  (equivalently  $r - q + 1 \leq k \leq t_2$ ), let  $j$  be the subscript with  $k = j + r - q$ . Then  $j = k - r + q$ . Thus  $1 \leq j \leq t_2 - r + q \leq q$  and hence  $F_{2,k}$  is a subset of  $F_j$  by (25). Therefore,

(26.5) there exists a subscript  $k$  with  $u_1 u_2 \in F_{2,k} \subseteq F_j$  iff  $t_1 + q + 1 \leq j \leq \min\{r, t_2 + q\}$  or  $1 \leq j \leq \max\{0, t_2 - r + q\}$ .

By the hypothesis of the present subcase,  $s_i + t_i + q = r$  for  $i = 1$  or  $2$ . If  $s_1 + t_1 + q = r$ , then  $r - s_1 + 1 = t_1 + q + 1$ . Clearly,  $r - s_2 \leq \min\{r, t_2 + q\}$ . Combining (26.4) (as  $F_{1,j} \subseteq F_j$ ) with (26.5), we see that  $F_j$  contains  $u_1 u_2$  iff  $t_1 + q + 1 \leq j \leq \min\{r, t_2 + q\}$  or  $1 \leq j \leq \max\{0, t_2 - r + q\}$ . If  $s_2 + t_2 + q = r$ , then  $r - s_2 = t_2 + q$ . Clearly,  $r - s_1 + 1 \leq t_1 + q + 1$ . It follows from (26.4) and (26.5) that  $F_j$  contains  $u_1 u_2$  iff  $r - s_1 + 1 \leq j \leq r - s_2$ . Thus (26) holds.

By (1), we have  $p \geq \max\{s_1 - s_2, t_2 - t_1\}$ . In view of (24)-(26), we obtain

(27) each arc  $e$  of  $T$  is contained in at most  $w(e)$  members of  $\mathcal{F} := \{F_1, F_2, \dots, F_r\}$ .

Let us show that

(28) each  $F_j$ , with  $1 \leq j \leq r$ , is an FAS of  $T$ .

For this purpose, let  $C$  be an arbitrary cycle in  $T$ . Clearly,  $F_j$  intersects  $C$  if  $C$  is a cycle of  $T_1$  or a cycle of  $T_2$ . So we assume that  $C$  is not fully contained in  $T_i$  for  $i = 1, 2$ .

Suppose  $C$  contains an arc  $ab$  with  $a \in V(A_1 \setminus v_1)$  and  $b \notin V(A_1 \cup B_1)$ . From the structural description, we see that  $C$  passes through  $v_1$  and  $C[v_1, a]$  is fully contained in  $A_1$ . Let  $C'$  be the cycle arising from  $C$  by replacing  $C[v_1, b]$  with  $v_1 b$ , and let  $F_{2,k}$  be the member of  $\mathcal{F}_2$  contained in  $F_j$ . Then  $C'$  is fully contained in  $T_2$  and intersects  $F_{2,k}$ . If  $F_{2,k}$  intersects  $C'[b, v_1] = C[b, v_1]$ , then  $F_j$  intersects  $C$ . So we assume that  $F_{2,k}$  contains  $v_1 b$  and hence  $w(v_1 b) > 0$ . It follows from (22) that  $q \geq w(v_1 b) > 0$ . By (24) and (25), we get  $k = j + r - q$  and  $F_{1,j}$  contains the

blue arc  $v_1b$  as well. By (1), we obtain  $w(C[v_1, b]) \geq w(v_1b)$ . From the construction of  $\mathcal{F}_1$  using breadth-first search, we see that  $F_{1,j}$  intersects  $C[v_1, b]$ . Thus  $F_j$  intersects  $C$ .

So we assume that  $C$  contains no arc  $ab$  with  $a \in V(A_1 \setminus v_1)$  and  $b \notin V(A_1 \cup B_1)$ . Consider the situation when  $C$  contains both  $v_2u_1v_1$  and  $au_2b$  as segments, where  $a \in V(A_1 \setminus v_1)$  and  $b \notin V(A_1 \cup B_1)$ . Note that  $C[v_1, a]$  is fully contained in  $A_1$ . Let  $C'$  be obtained from  $C$  by replacing  $C[v_2, u_2]$  with  $v_2u_2$ . Then  $C'$  is fully contained in  $T_2$ . So  $F_j$  intersects  $C'$ . If  $v_2u_2 \notin F_j$ , then  $F_j$  intersects  $C'[u_2, v_2]$  and hence  $C$ ; otherwise,  $v_2u_2 \in F_j$ , so  $q+1 \leq j \leq \min\{r, q+t_2\}$  or  $1 \leq j \leq \max\{0, t_2-r+q\}$  by (26). If  $q+1 \leq j \leq r$  then, by (26),  $F_j$  contains  $v_2u_1$  or  $u_1v_1$ , because  $q+t_1 \geq r-s_1$ . So  $F_j$  intersects  $C$ . If  $1 \leq j \leq t_2-r+q$  then  $q+t_2 > r$  and hence  $s_1+t_1+q=r$  by the hypothesis of Subcase 2.1. By (1), we have  $t_1+s_1+(r-s_2) \geq t_2$  (when  $s_2 > 0$ , arc  $u_2v_1$  is contained in a cycle  $Q$  of  $T_1$  having weight  $r$  with respect to  $w$  by (6). Consider the path  $v_2u_1v_1Q[v_1, u_2]$ , which has weight  $t_1+s_1+(r-s_2)$ ). It follows that  $t_2-r+q \leq r-s_2$ . Thus  $1 \leq j \leq r-s_2$ . So  $F_{1,j}$  intersects  $C[v_1, u_2]$  by (9) and hence  $F_j$  intersects  $C$  by (25).

Notice that  $u_1u_2$  plays no role in the above proof. So the same argument (simply interchanging the subscripts 1 and 2, whenever appropriate) implies that  $F_j$  also intersects  $C$  if  $C$  contains both  $v_2u_2v_1$  and  $au_1b$  as segments, where  $a \in V(A_1 \setminus v_1)$  and  $b \notin V(A_1 \cup B_1)$ . This proves (28).

Combining (27) with (28), we conclude that  $\mathcal{F}$  is a  $w$ -FAS packing of  $T$  having size  $r$ . From (9) and (25), it is clear that  $\mathcal{F} \cap E^*$  is obtained by first performing breadth-first search for  $|\mathcal{F} \cap E^*|$  steps in  $T^*$  from  $v_1$  and then eliminating triangles in  $A_1 \setminus v_1$ .

**Subcase 2.2.**  $s_i+t_i+q > r$  for  $i=1, 2$ .

Recall that each arc  $e$  with  $w(e) > 0$  is contained in a minimum cycle of  $(T, w)$ . By (21), we obtain

(29)  $q=0$ . So  $s_i+t_i > r$  for  $i=1, 2$  and hence  $s_1, s_2, t_1$  and  $t_2$  are all positive.

In view of (21) and (29),  $R_1$  is contained in no cycle of  $T_i$  having weight  $r$  with respect to  $w$  for  $i=1, 2$ . It follows from (8) that

(30)  $w(ab)=0$  for any  $a \in V(A_1)$  and  $b \notin V(A_1 \cup B_1 \setminus v_2)$ .

For  $i=1, 2$ , let  $T'_i = (V'_i, E'_i)$  be obtained from  $T_i$  by deleting the vertex  $v_{3-i}$ , and let  $\mathcal{F}'_i = \{F'_{i,1}, F'_{i,2}, \dots, F'_{i,r}\}$ , where  $F'_{i,j}$  is the restriction of  $F_{i,j}$  to  $E'_i$  for  $1 \leq j \leq r$ . Observe that no arc is shared by a member of  $\mathcal{F}'_1$  and that of  $\mathcal{F}'_2$ , except  $u_1u_2$ . We shall produce a  $w$ -FAS packing  $\mathcal{F}$  of  $T$  having size  $r$  by gluing members of  $\mathcal{F}'_1$  together with those of  $\mathcal{F}'_2$ , along  $u_1u_2$  whenever possible. For this purpose, observe from (6), (9), (10), and (29) that

(31)  $F'_{1,j}$  contains  $u_1u_2$  iff  $r-s_1+1 \leq j \leq r-s_2$ , and  $F'_{2,k}$  contains  $u_1u_2$  iff  $t_1+1 \leq k \leq t_2$ ;

(32)  $F'_{1,j}$  contains  $u_iv_1$  iff  $r-s_i+1 \leq j \leq r$ , and no  $F'_{2,k}$  contains  $u_iv_1$  for  $i=1, 2$ ; and

(33) no  $F'_{1,j}$  contains  $v_2u_i$ , and  $F'_{2,k}$  contains  $v_2u_i$  iff  $1 \leq k \leq t_i$  for  $i=1, 2$ .

Let  $\{g, h\}$  be a permutation of  $\{1, 2\}$  with  $s_g+t_g \leq s_h+t_h$ . We first arrange  $F'_{1,1}, F'_{1,2}, \dots, F'_{1,r}$  on a circle  $O$  in clockwise order, and then arrange  $F'_{2,1}, F'_{2,2}, \dots, F'_{2,r}$  on  $O$  in the same order, such that members of  $\mathcal{F}'_1$  alternate with those of  $\mathcal{F}'_2$  in the following way:

- $F'_{2,t_g+1}$  follows  $F'_{1,r-s_g+1}$  immediately;
- $F'_{2,t_g+2}$  follows  $F'_{1,r-s_g+2}$  immediately;
- • • • •
- $F'_{2,t_g}$  follows  $F'_{1,r-s_g}$  immediately,

where the subscripts are taken modulo  $r$ . In particular,  $F'_{i,0} = F'_{i,r}$  for  $i=1, 2$ .

For  $1 \leq j \leq r$ , let  $\pi(j)$  denote the subscript such that  $F'_{2,\pi(j)}$  follows  $F'_{1,j}$  immediately on  $O$ , and define  $F_j = F'_{1,j} \cup F'_{2,\pi(j)}$ . Observe that

$$(34) \quad \pi(j) = \begin{cases} (s_g + t_g - r) + j & \text{if } 1 \leq j \leq 2r - (s_g + t_g), \\ (s_g + t_g - 2r) + j & \text{if } 2r - (s_g + t_g) + 1 \leq j \leq r, \end{cases} \quad \text{which implies } \pi(r - s_g) = t_g \text{ if } s_g < r \text{ and } \pi(r - s_h) \leq t_h \text{ if } s_h < r \text{ (the first line of } \pi(j) \text{ applies now).}$$

(35) Each  $F_j$  for  $1 \leq j \leq r$  intersects each of the three paths  $v_2u_1v_1$ ,  $v_2u_2v_1$ , and  $v_2u_1u_2v_1$ .

To justify this, imagine that circle  $O$  has  $r$  positions,  $1, 2, \dots, r$ , in clockwise order, such that each position  $i$  is occupied by both  $F'_{1,i}$  and  $F'_{2,\pi(i)}$ . By (29), we have  $s_g + t_g > r$ . From the arrangements of  $F_{i,j}$ 's on  $O$ , it follows immediately that

(35.1) circle  $O$  is covered by  $F'_{1,r-s_g+1}, F'_{1,r-s_g+2}, \dots, F'_{1,r}, F'_{2,1}, F'_{2,2}, \dots, F'_{2,t_g}$ ; that is, each position of  $O$  is occupied by at least one of these sets.

(35.2) Circle  $O$  is also covered by  $F'_{1,r-s_h+1}, F'_{1,r-s_h+2}, \dots, F'_{1,r}, F'_{2,1}, F'_{2,2}, \dots, F'_{2,t_h}$ .

The statement holds trivially if  $s_h = r$ . So we assume that  $s_h < r$ . From (34) and (29) we deduce that  $\pi(1) = (s_g + t_g - r) + 1 \geq 2$  and  $\pi(r - s_h) \leq t_h$ . Hence  $\{F'_{2,\pi(1)}, F'_{2,\pi(2)}, \dots, F'_{2,\pi(r-s_h)}\} \subseteq \{F'_{2,1}, F'_{2,2}, \dots, F'_{2,t_h}\}$ , this proves (35.2).

Similarly, we can check that  $\{F'_{2,\pi(1)}, F'_{2,\pi(2)}, \dots, F'_{2,\pi(r-s_2)}\} \subseteq \{F'_{2,1}, F'_{2,2}, \dots, F'_{2,t_1}, F'_{2,t_1+1}, F'_{2,t_1+2}, \dots, F'_{2,t_2}\}$ , where  $F'_{2,t_1+1}, F'_{2,t_1+2}, \dots, F'_{2,t_2}$  appear only when  $t_1 < t_2$ . Thus

(35.3) circle  $O$  is, moreover, covered by  $F'_{1,r-s_2+1}, F'_{1,r-s_2+2}, \dots, F'_{1,r}, F'_{2,1}, F'_{2,2}, \dots, F'_{2,t_1}, F'_{2,t_1+1}, F'_{2,t_1+2}, \dots, F'_{2,t_2}$ .

Combining (31)-(33) and (35.1)-(35.3), we conclude that each  $F_j$  for  $1 \leq j \leq r$  intersects each of the three paths  $v_2u_1v_1$ ,  $v_2u_2v_1$ , and  $v_2u_1u_2v_1$ .

(36)  $u_1u_2$  is contained in at most  $w(u_1u_2)$  members of the family  $\mathcal{F} := \{F_1, F_2, \dots, F_r\}$ .

Since  $\mathcal{F}'_i$  ( $i = 1, 2$ ) is obtained by restricting the  $w_i$ -packing  $\mathcal{F}_i$  to  $E'_i$ , the construction of  $\mathcal{F}$  and (31) allow us to assume that  $s_2 + 1 \leq s_1$  and  $t_1 + 1 \leq t_2$ . By (1) with respect to  $w_1$  and  $w_2$  respectively, we obtain  $s_1 - s_2 \leq w_1(u_1u_2)$  and  $t_2 - t_1 \leq w_2(u_1u_2)$ . Hence  $\max\{s_1 - s_2, t_2 - t_1\} \leq \max\{w_1(u_1u_2), w_2(u_1u_2)\} \leq w(u_1u_2)$ . When  $g = 1$ , it is instant from the construction of  $\mathcal{F}$  that exactly  $\max\{s_1 - s_2, t_2 - t_1\}$  members of  $\mathcal{F}$  contain  $u_1u_2$ . When  $g = 2$ , since  $r - s_1 + 1 \geq 1$  and  $t_1 \leq t_2 - 1$ , it follows from (34) (the first line) that  $\pi(r - s_1 + 1) \leq t_1 + 1$ . Thus  $\pi(r - s_1 + 1) \leq t_2 = \pi(r - s_2)$ , which implies that exactly  $s_1 - s_2$  members of  $\mathcal{F}$  contain  $u_1u_2$ . Therefore (36) holds in either case.

(37) Each  $F_j$ , with  $1 \leq j \leq r$ , is an FAS of  $T$ .

To see this, let  $C$  be an arbitrary cycle in  $T$ . Clearly,  $F_j$  intersects  $C$  if  $C$  is a cycle of  $T'_1$  or a cycle of  $T'_2$ . So we assume that  $C$  is not fully contained in  $T'_i$  for  $i = 1, 2$ . From the structural description of  $T$ , we deduce that  $C$  contains one of the three paths  $v_2u_1v_1$ ,  $v_2u_2v_1$ , and  $v_2u_1u_2v_1$  as a segment. Therefore  $F_j$  intersects  $C$  by (35), as desired.

Since no arc is shared by a member of  $\mathcal{F}'_1$  and that of  $\mathcal{F}'_2$ , except  $u_1u_2$ , the family  $\mathcal{F} = \{F_1, F_2, \dots, F_r\}$  is a  $w$ -FAS packing of  $T$  having size  $r$  by (36) and (37). From (9) and (30), it is clear that  $\mathcal{F} \cap E^*$  is obtained by first performing breadth-first search for  $|\mathcal{F} \cap E^*|$  steps in  $T^*$  from  $v_1$  and then eliminating triangles in  $A_1 \setminus v_1$ . This completes the proof of Theorem 3.3.  $\blacksquare$

## 4 Computer-assisted Proof

In the preceding section we have established the desired minimax relation for all Möbius-free strong tournaments other than  $F_1$  and  $G_1$ , thereby finishing the main body of the proof of Theorem 1.2. In this section we present a computer-assisted proof for  $G_1$ .

**Lemma 4.1.** *Tournament  $G_1$  is FAS Mengerian.*

In Schrijver [27] there is a characterization (Corollary 22.13d) of TDI system of the form  $Ax \leq b, x \geq \mathbf{0}$ , where  $A$  is a nonnegative integral matrix. The same argument yields the following result.

**Lemma 4.2.** *Let  $A$  be a nonnegative integral matrix with no zero rows, and let  $b$  be a rational vector. Then the system  $Ax \geq b, x \geq \mathbf{0}$  is TDI iff for each  $\{0,1\}$ -vector  $y$ , there exists an integral vector  $z \geq \mathbf{0}$  with  $z^T A \leq \lceil y^T A/2 \rceil$  and  $2z^T b \geq y^T b$ .*

To prove Lemma 4.1, let  $A$  be the minimal FAS-arc incidence matrix of  $G_1$ . Clearly,  $G_1$  is FAS Mengerian iff  $Ax \geq \mathbf{1}, x \geq \mathbf{0}$  is a TDI system. We shall demonstrate that the dimension of  $A$  is  $41 \times 15$ . Since it is beyond the capacity of our computer to exhaust all possible  $2^{41}$  cases addressed in Lemma 4.2, we have to derive a refinement of this lemma to fulfill our need.

Suppose the dimension of  $A$  in Lemma 4.2 is  $m \times n$ . Let  $\prec$  denote the lexicographical order defined over the set of all  $m$ -dimensional  $\{0,1\}$ -vectors; that is,  $u \prec v$  if there exists a subscript  $j$ , with  $1 \leq j \leq m$ , such that  $u_i = v_i$  for all  $1 \leq i < j$  and  $u_j < v_j$ .

**Lemma 4.3.** *Let  $A$  be a nonnegative integral matrix with no zero rows. Let  $V$  and  $W$  be two sets of  $\{0,1\}$ -vectors such that for each  $v \in V$ , there exists  $w \in W$  satisfying  $v \prec w, v^T \mathbf{1} = w^T \mathbf{1}$ , and  $w^T A \leq v^T A$ . Let  $U$  consist of all  $\{0,1\}$ -vectors  $u$  such that  $u^T \mathbf{1}$  is odd and  $u \not\geq v$  for each  $v \in V$ . Then the system  $Ax \geq \mathbf{1}, x \geq \mathbf{0}$  is TDI iff for each  $y \in U$ , there exists an integral vector  $z \geq \mathbf{0}$  with  $z^T A \leq \lceil y^T A/2 \rceil$  and  $2z^T \mathbf{1} \geq y^T \mathbf{1}$ .*

**Proof.** The “only if” part follows instantly from Lemma 4.2.

To establish the “if” part, it suffices to find a desired  $z$  for every  $\{0,1\}$ -vector  $y$  as described in Lemma 4.2. Suppose on the contrary that such  $z$  does not exist for some  $y$ . We choose such a counterexample  $y$  with the property that

- (1)  $y^T \mathbf{1}$  is as small as possible, and
- (2) subject to (1), the lexicographical order of  $y$  is as high as possible.

Note that  $y \notin U$ , and thus either  $y^T \mathbf{1}$  is even or  $y \geq v$  for some  $v \in V$ .

We first assume that  $y^T \mathbf{1}$  is even. Now  $y \neq \mathbf{0}$ , for otherwise  $z = \mathbf{0}$  would satisfy the requirements. Thus there exists a unit  $\{0,1\}$ -vector  $e \leq y$ . Since  $(y - e)^T \mathbf{1} = y^T \mathbf{1} - 1$ , condition (1) guarantees the existence of an integral vector  $z \geq \mathbf{0}$  satisfying  $z^T A \leq \lceil (y - e)^T A/2 \rceil$  and  $2z^T \mathbf{1} \geq (y - e)^T \mathbf{1}$ , which clearly imply  $z^T A \leq \lceil y^T A/2 \rceil$  and  $2z^T \mathbf{1} \geq y^T \mathbf{1}$ , a contradiction.

Next, we assume that  $y \geq v$  for some  $v \in V$ . By hypothesis, there exists  $w \in W$  such that  $v \prec w, v^T \mathbf{1} = w^T \mathbf{1}$ , and  $w^T A \leq v^T A$ . Observe that  $y - v + w$  can be expressed as  $\alpha + 2\beta$  for some  $\{0,1\}$ -vectors  $\alpha$  and  $\beta$ . We proceed by considering two subcases.

Suppose  $\beta \neq \mathbf{0}$ . Then  $\alpha^T \mathbf{1} = y^T \mathbf{1} - v^T \mathbf{1} + w^T \mathbf{1} - 2\beta^T \mathbf{1} = y^T \mathbf{1} - 2\beta^T \mathbf{1} < y^T \mathbf{1}$ . By (1), there exists an integral vector  $\gamma \geq \mathbf{0}$  satisfying  $\gamma^T A \leq \lceil \alpha^T A/2 \rceil$  and  $2\gamma^T \mathbf{1} \geq \alpha^T \mathbf{1}$ . Set  $z = \gamma + \beta$ . Then  $z^T A = \gamma^T A + \beta^T A \leq \lceil \alpha^T A/2 \rceil + \beta^T A = \lceil (\alpha + 2\beta)^T A/2 \rceil = \lceil (y - v + w)^T A/2 \rceil \leq \lceil y^T A/2 \rceil$ .

Similarly,  $2z^T \mathbf{1} = 2\gamma^T \mathbf{1} + 2\beta^T \mathbf{1} \geq \alpha^T \mathbf{1} + 2\beta^T \mathbf{1} = (y - v + w)^T \mathbf{1} = y^T \mathbf{1}$ , which is impossible as  $y$  is a counterexample.

Suppose  $\beta = \mathbf{0}$ . Then  $\alpha^T \mathbf{1} = (y - v + w)^T \mathbf{1} = y^T \mathbf{1}$ . Since  $v \prec w$ , we have  $y \prec \alpha$ , which implies, from (2), the existence of an integral vector  $z \geq \mathbf{0}$  such that  $z^T A \leq \lceil \alpha^T A / 2 \rceil$  and  $2z^T \mathbf{1} \geq \alpha^T \mathbf{1}$ . Consequently,  $z^T A \leq \lceil (y - v + w)^T A / 2 \rceil \leq \lceil y^T A / 2 \rceil$  and  $2z^T \mathbf{1} \geq (y - v + w)^T \mathbf{1} = y^T \mathbf{1}$ , again a contradiction. ■

As we shall see, Lemma 4.3 can help eliminate many cases involved in our analysis.

**Proof of Lemma 4.1.** Tournament  $G_1$  is as shown in Figure 5. For simplicity, we relabel each vertex  $v_i$  as  $i$  for  $1 \leq i \leq 6$ . Thus the vertex set of  $G_1$  is  $V_1 = \{1, 2, 3, 4, 5, 6\}$  and arc set is  $E_1 = \{12, 23, 34, 45, 51, 13, 35, 52, 24, 41, 16, 26, 63, 64, 65\}$  whose members are denoted by  $a, b, c, d, e, f, g, h, i, j, k, l, m, n, o$ , respectively (so  $a = 12, b = 23, c = 34$  and so on).

**Claim 1.** Let  $\mathcal{F}$  be the family of all minimal feedback arc sets of  $G_1$ . Then  $|\mathcal{F}| = 41$  and  $\mathcal{F} = \{ehj, afhk, dgjo, acehk, acehn, acghk, bdejl, beijl, bfikl, cehin, cgikl, cgin, degjl, dgjkl, abdfkl, acdgkl, acdgko, acdgno, acghno, adfgkl, adfgko, adghjk, aefhmn, afhmn, bceikl, bceiln, bdejmo, bdfjkl, bdfjmo, bfirmno, cegijl, cegiln, cehikl, abdfkmo, abdfmno, adfgmno, adfhjmo, bceimno, befhimn, befilmn, beijmno\}$ , where, for instance,  $ehj$  stands for the minimal FAS consisting of arcs  $e, h$  and  $j$ .

To justify this, we first list all subsets of  $E_1$  in nondecreasing order of cardinality. For each term  $F$  on the list, from the first to the last, we check if  $G_1 \setminus F$  is acyclic and if  $F$  contains a feedback arc set we have already found. If  $F$  is a feedback arc set and it does not contain any earlier ones, then  $F$  is a minimal feedback arc set and we put it in  $\mathcal{F}$ . When the process is finished, we end up with 41 minimal feedback arc sets as shown above. This step was carried out by using computer.

Let  $A$  be the minimal FAS-arc incidence matrix of  $G_1$ , such that the  $i$ th row of  $A$  corresponds to the  $i$ th member of  $\mathcal{F}$  displayed in Claim 1. We shall use Lemma 4.3 to verify that the system  $Ax \geq \mathbf{1}, x \geq \mathbf{0}$  is TDI. To this end, let  $S_V$  and  $S_W$  be two families of 2-subsets of  $\{1, 2, \dots, 41\}$  as defined below (the subset  $\{i, j\}$  is written as  $i-j$ ):

$S_V = \{2-7, 2-8, 2-13, 2-27, 2-31, 2-41, 3-4, 3-5, 3-10, 3-23, 3-33, 3-39, 4-8, 4-9, 4-13, 4-14, 4-20, 4-21, 4-22, 4-23, 4-24, 4-28, 4-29, 4-30, 4-31, 4-36, 4-37, 4-39, 4-40, 4-41, 5-8, 5-9, 5-14, 5-21, 5-22, 5-28, 5-29, 5-31, 5-37, 5-41, 6-7, 6-8, 6-9, 6-13, 6-15, 6-20, 6-21, 6-23, 6-25, 6-26, 6-27, 6-28, 6-29, 6-30, 6-31, 6-32, 6-33, 6-34, 6-35, 6-36, 6-37, 6-38, 6-39, 6-40, 6-41, 7-10, 7-12, 7-19, 7-21, 7-22, 7-24, 7-33, 7-37, 7-39, 8-10, 8-15, 8-16, 8-17, 8-18, 8-19, 8-21, 8-22, 8-23, 8-24, 8-28, 8-33, 8-35, 8-36, 8-37, 8-39, 9-19, 9-22, 9-23, 9-24, 9-31, 9-37, 9-41, 10-13, 10-14, 10-15, 10-16, 10-17, 10-18, 10-19, 10-20, 10-21, 10-22, 10-27, 10-28, 10-29, 10-31, 10-34, 10-35, 10-37, 10-41, 11-15, 11-22, 11-23, 11-24, 11-27, 11-29, 11-34, 11-35, 11-37, 11-39, 11-41, 12-13, 12-14, 12-15, 12-16, 12-22, 12-27, 12-28, 12-29, 12-33, 12-37, 12-39, 13-17, 13-19, 13-21, 13-22, 13-24, 13-25, 13-27, 13-29, 13-30, 13-33, 13-34, 13-37, 13-38, 13-39, 13-41, 14-17, 14-18, 14-19, 14-21, 14-23, 14-24, 14-25, 14-26, 14-27, 14-30, 14-31, 14-32, 14-33, 14-34, 14-35, 14-36, 14-37, 14-38, 14-39, 14-41, 15-19, 15-22, 15-24, 15-30, 15-31, 15-33, 15-37, 15-39, 15-41, 16-19, 16-22, 16-24, 16-28, 16-29, 16-30, 16-31, 16-33, 16-37, 16-39, 16-40, 16-41, 17-19, 17-22, 17-23, 17-24, 17-28, 17-29, 17-30, 17-31, 17-32, 17-33, 17-37, 17-39, 17-40, 17-41, 18-22, 18-28, 18-29, 18-30, 18-31, 18-33, 18-37,$

18-39, 18-41, 19-20, 19-21, 19-22, 19-25, 19-26, 19-27, 19-28, 19-29, 19-31, 19-32, 19-33, 19-34, 19-35, 19-36, 19-37, 19-38, 19-39, 19-40, 19-41, 20-22, 20-24, 20-25, 20-26, 20-27, 20-28, 20-29, 20-30, 20-31, 20-33, 20-37, 20-38, 20-39, 20-41, 21-22, 21-23, 21-24, 21-25, 21-26, 21-27, 21-28, 21-29, 21-31, 21-33, 21-37, 21-38, 21-39, 21-40, 21-41, 22-23, 22-24, 22-25, 22-26, 22-27, 22-28, 22-29, 22-30, 22-31, 22-32, 22-33, 22-34, 22-35, 22-36, 22-37, 22-38, 22-39, 22-40, 22-41, 23-25, 23-27, 23-28, 23-29, 23-31, 23-33, 23-34, 23-37, 23-38, 23-41, 24-25, 24-26, 24-27, 24-28, 24-31, 24-33, 24-34, 24-38, 24-39, 24-40, 24-41, 25-28, 25-31, 25-35, 25-36, 25-37, 25-39, 25-41, 26-28, 26-29, 26-31, 26-33, 26-34, 26-36, 26-37, 26-39, 26-41, 27-31, 27-32, 27-33, 27-36, 27-37, 27-39, 27-40, 28-31, 28-32, 28-33, 28-34, 28-35, 28-36, 28-37, 28-38, 28-39, 28-40, 28-41, 29-31, 29-32, 29-33, 29-36, 29-38, 29-39, 29-40, 29-41, 30-31, 30-33, 30-37, 31-33, 31-34, 31-35, 31-36, 31-37, 31-38, 31-39, 31-40, 31-41, 32-33, 32-34, 32-35, 32-37, 32-39, 32-41, 33-34, 33-35, 33-36, 33-37, 33-38, 33-39, 33-40, 33-41, 34-37, 34-39, 34-40, 34-41, 35-37, 35-39, 35-41, 36-37, 36-38, 36-39, 36-41, 37-38, 37-39, 37-40, 37-41, 38-39, 39-41, 40-41} and

$S_W = \{1-2, 1-3, 1-6, 1-9, 1-11, 1-12, 1-14, 1-15, 1-16, 1-17, 1-18, 1-20, 1-24, 1-25, 1-26, 1-29, 1-30, 1-32, 1-34, 1-35, 1-36, 1-38, 1-40, 2-3, 2-5, 2-10, 2-11, 2-12, 2-14, 2-16, 2-17, 2-18, 2-25, 2-26, 2-29, 2-30, 2-32, 2-35, 2-36, 2-38, 2-40, 3-6, 3-7, 3-8, 3-9, 3-11, 3-15, 3-16, 3-20, 3-24, 3-25, 3-26, 3-30, 3-32, 3-34, 3-35, 3-38, 3-40, 4-11, 4-12, 5-11, 5-12, 5-30, 6-14, 6-18, 7-9, 7-11, 7-30, 8-11, 8-12, 8-30, 8-32, 8-38, 9-10, 9-16, 9-17, 9-18, 9-35, 9-36, 10-11, 10-25, 10-30, 10-40, 11-13, 11-18, 12-34, 12-35, 14-15, 15-29, 15-32, 15-38, 18-24, 18-40, 23-30, 24-29, 27-30\}$ .

Notice that  $|S_V| = 390$  and  $|S_W| = 96$ ; these  $S_V$  and  $S_W$  will yield  $V$  and  $W$  as described in Lemma 4.3. The choices for  $S_V$  and  $S_W$  are not unique. We obtained our  $S_V$  and  $S_W$  by trail and error. In the search process we restricted our attention to 2-subsets. It is possible to choose larger sets  $S_V$  and  $S_W$ , which would cause  $\Gamma$  (to be defined in Claim 3) to contain fewer stable sets.

**Claim 2.** Let  $V$  and  $W$  be the sets of characteristic vectors (with length 41) of members of  $S_V$  and  $S_W$ , respectively. Then  $V$  and  $W$  satisfy the conditions described in Lemma 4.3.

To justify this, for each of the 390 vectors  $v \in V$  and each of the 96 vectors  $w \in W$ , we test if  $v \prec w$  and  $w^T A \leq v^T A$  hold simultaneously (note that  $v^T \mathbf{1} = w^T \mathbf{1}$  is always true). Using a computer we have confirmed that, for every  $v \in V$  indeed there exists  $w \in W$  such that  $v \prec w$  and  $w^T A \leq v^T A$  are both true.

**Claim 3.** Let  $\Gamma$  be the graph with vertex set  $\{1, 2, \dots, 41\}$  such that  $i, j \in \{1, 2, \dots, 41\}$  are adjacent iff  $i-j$  is a member of  $S_V$ . Then  $\Gamma$  has exactly 41022 odd stable sets.

Mathematica has a function *FindClique*, which can be used to generate all 219 maximal stable sets of  $\Gamma$ . We also independently implemented the Bron-Kerbosch algorithm (see [https://en.wikipedia.org/wiki/Bron-Kerbosch\\_algorithm](https://en.wikipedia.org/wiki/Bron-Kerbosch_algorithm)) and obtained the same result. These maximal stable sets give rise to all 82044 stable sets, and exactly half of which are odd.

**Claim 4.** System  $Ax \geq \mathbf{1}$ ,  $x \geq \mathbf{0}$  is TDI.

To justify this, let us choose  $V$  and  $W$  as in Claim 2. It is then clear that  $U$  (defined in Lemma 4.3) consists of exactly characteristic vectors of odd stable sets of  $\Gamma$ . By Claim 3,  $|U| = 41022$ . For each  $y \in U$ , we define  $c = \lceil y^T A / 2 \rceil$  and solve  $\max\{z^T \mathbf{1} : z^T A \leq c^T, z \geq \mathbf{0} \text{ and integral}\}$  using *LinearProgramming* of Mathematica. For each optimal solution  $z$  obtained, we verify that  $2z^T \mathbf{1} \geq y^T \mathbf{1}$ . We also verify that  $z$  is an integral vector satisfying  $z^T A \leq c^T$ . Our

computational results indicate that indeed that is the case. After completing this process for all 41022 vectors in  $U$ , we conclude from Lemma 4.3 that Claim 4 is true. ■

We can finally establish the equivalence of three statements described in Theorem 1.2, thereby obtaining a complete characterization of all FAS ideal and Mengerian tournaments.

**Proof of Theorem 1.2.** Implication  $(iii) \Rightarrow (ii)$  holds, because total-dual integrality implies primal integrality (see Edmonds-Giles theorem [15] stated in Section 1). It was proved by Lehman [22] that a clutter is ideal iff its blocker is ideal, which implies that a tournament is cycle ideal iff it is FAS ideal. Therefore the equivalence of (i) and (ii) in Theorem 1.1 yields implication  $(ii) \Rightarrow (i)$ . It remains to prove implication  $(i) \Rightarrow (iii)$ . Clearly, we may assume that  $T$  is strong. Since  $F_1$  arises from  $G_1$  by deleting vertex  $v_6$  (see the labeling in Figure 5), from Lemma 4.1 we deduce that  $F_1$  is also FAS Mengerian. So we may assume that  $T \notin \{F_1, G_1\}$ . From Theorem 3.1 we thus conclude that  $T$  is FAS Mengerian. ■

## References

- [1] N. Alon, Ranking tournaments, *SIAM J. Discrete Math.* **20** (2006), 137-142.
- [2] D. Applegate, W. Cook, and S. McCormick, Integral infeasibility and testing total dual integrality, *Oper. Res. Lett.* **10** (1991), 37-41.
- [3] F. Barahona, J. Fonlupt, and A. Mahjoub, Compositions of graphs and polyhedra IV: Acyclic spanning subgraphs, *SIAM J. Discrete Math.* **7** (1994), 390-402.
- [4] F. Barahona and A. Mahjoub, Composition in the acyclic subdigraph polytope, Report No. 85371-OR, Institut für Ökonometrie und Operations Research, Universität Bonn, 1985.
- [5] M. Cai, X. Deng, and W. Zang, An approximation algorithm for feedback vertex sets in tournaments, *SIAM J. Comput.* **30** (2001), 1993-2007.
- [6] X. Chen, G. Ding, X. Hu, and W. Zang, A min-max relation on packing feedback vertex sets, *Math. Oper. Res.* **31** (2006), 777-788.
- [7] X. Chen, G. Ding, W. Zang, and Q. Zhao, Ranking tournaments with no errors I: Structural description, *J. Combin. Theory Ser. B* **141** (2020), 264-294.
- [8] X. Chen, G. Ding, W. Zang, and Q. Zhao, Ranking tournaments with no errors II: Minimax relation, *J. Combin. Theory Ser. B* **142** (2020), 244-275.
- [9] X. Chen, X. Hu, and W. Zang, A min-max theorem on tournaments, *SIAM J. Comput.* **37** (2007), 923-937.
- [10] P. Charbit, P. Thomassé, and A. Yeo, The minimum feedback arc set problem is *NP*-hard for tournaments, *Combin. Probab. Comput.* **16** (2007), 1-4.
- [11] G. Cornuéjols and B. Guenin, On dijoins, *Discrete Math.* **243** (2002), 213-216.
- [12] M. Chudnovsky, K. Edwards, R. Kim, A. Scott, and P. Seymour, Disjoint dijoins, *J. Combin. Theory Ser. B* **120** (2016), 18-35.
- [13] G. Ding, Z. Xu, and W. Zang, Packing cycles in graphs, II, *J. Combin. Theory Ser. B* **87** (2003), 244-253.

- [14] G. Ding and W. Zang, Packing cycles in graphs, *J. Combin. Theory Ser. B* **86** (2002), 381-407.
- [15] J. Edmonds and R. Giles, A min-max relation for submodular functions on graphs, in: *Ann. Discrete Math.* **1**, North-Holland, Amsterdam, 1977, pp. 185-204.
- [16] P. Feofiloff and D. Younger, Directed cut transversal packing for source-sink connected graphs, *Combinatorica* **7** (1987), 255-263.
- [17] M. Grötschel, L. Lovász, and A. Schrijver, The ellipsoid method and its consequences in combinatorial optimization, *Combinatorica* **1** (1981), 169-197.
- [18] B. Guenin, Circuit Mengerian directed graphs, in: *Integer Programming and Combinatorial Optimization* (Utrecht, 2001), Lecture Notes in Comput. Sci. 2081, pp. 185-195.
- [19] B. Guenin and R. Thomas, Packing directed circuits exactly, *Combinatorica* **31** (2011), 397-421.
- [20] M. Jünger, *Polyhedral Combinatorics and the Acyclic Subdigraph Problem*, Heldermann Verlag, Berlin, 1985.
- [21] O. Lee and Y. Wakabayashi, Note on a min-max conjecture of Woodall, *J. Graph Theory* **38** (2001), 36-41.
- [22] A. Lehman, On the length-width inequality, *Math. Programming* **17** (1979), 403-417.
- [23] C. Lucchesi and D. Younger, A minimax theorem for directed graphs, *J. London Math. Soc.* **17** (1978), 369-374.
- [24] A. Mészáros, A note on disjoint dijoins, *Combinatorica* **38** (2018), 1485–1488.
- [25] A. Schrijver, A counterexample to a conjecture of Edmonds and Giles, *Discrete Math.* **32** (1980), 213-214.
- [26] A. Schrijver, Min-max relations for directed graphs, *Ann. Discrete Math.* **16** (1982), 261-280.
- [27] A. Schrijver, *Theory of Linear and Integer Programming*, John Wiley & Sons, New York, 1986.
- [28] P. Seymour, Packing circuits in eulerian digraphs, *Combinatorica* **16** (1996), 223-231.
- [29] A. Williams and B. Guenin, Advances in packing directed joins, *Electron. Notes Discrete Math.* **19** (2005), 249-255.
- [30] D.R. Woodall, Menger and König systems, *Lecture Notes in Math.* **642** (1978), 620-635.