Weighted Composition Operators between Different Bloch-type Spaces in Polydiscs

Zehua Zhou
Department of Mathematics, Tianjin University
Tianjin 300072, P. R. China
LiuHui Center for Applied Mathematics
Nankai University and Tianjin University
Tianjin 300072, P. R. China
E-mail: zehuazhou@hotmail.com

1

Abstract

Let $\phi(z) = (\phi_1(z), \dots, \phi_n(z))$ be a holomorphic self-map of \mathbb{D}^n and $\psi(z)$ a holomorphic function on \mathbb{D}^n , where \mathbb{D}^n is the unit polydiscs of \mathbb{C}^n . Let p > 0, q > 0, this paper gives some necessary and sufficient conditions for the weighted composition operator $W_{\psi,\phi}$ induced by ψ and ϕ to be bounded and compact between Bloch-type space $\beta_p(\mathbb{D}^n)$ and $\beta_q(\mathbb{D}^n)$.

 ${\bf Keywords}~$ Bloch-type space, Weighted composition operator, Composition operator.

2000 Mathematics Subject Classification 47B38, 47B33, 32A37, 32A30

¹Supported in part by the National Natural Science Foundation of China (Grand No.10001030), and LiuHui Center for Applied Mathematics, Nankai University and Tianjin University