# CONSTRUCTION OF SIMPLE QUOTIENTS OF BERNSTEIN-ZELEVINSKY DERIVATIVES AND HIGHEST DERIVATIVE MULTISEGMENTS II: MINIMAL SEQUENCES

#### KEI YUEN CHAN

ABSTRACT. Let F be a non-Archimedean local field. For any irreducible smooth representation  $\pi$  of  $\operatorname{GL}_n(F)$  and a multisegment  $\mathfrak{m}$ , we have an operation  $D_{\mathfrak{m}}(\pi)$  to construct a simple quotient  $\tau$  of a Bernstein-Zelevinsky derivative of  $\pi$ . This article continues the previous one to study the following poset

#### $\mathcal{S}(\pi,\tau) := \left\{ \mathfrak{n} : D_{\mathfrak{n}}(\pi) \cong \tau \right\},\,$

where **n** runs for all the multisegments. Here the partial ordering on  $S(\pi, \tau)$  comes from the Zelevinsky ordering. We show that the poset has a unique minimal multisegment. Along the way, we introduce two new ingredients: fine chain orderings and local minimizability.

### 1. INTRODUCTION

The Bernstein-Zelevinsky (BZ) derivative is a classical tool in studying the representation theory of p-adic groups, which is originally introduced in [Ze80]. In [Ch25], we transfer several problems of the study of simple quotients into sequences of derivatives of essentially square-integrable representations, that is to obtain some specific information of Jacquet modules. In this article, we further study the properties of sequences of derivatives, and some more in the sequels [Ch24+e].

One classical viewpoint of Jacquet modules (and its adjoint functor- parabolic induction) is the Hopf algebra approach, see e.g. [HM08]. Such study is very successful and useful, but it is usually carried in Grothendieck group level. The semisimplification of Jacquet modules sometimes has nicer formula e.g. [Ze80] and [Ma13, MT15] (for other classical groups), but it could be also hard for some special cases, see e.g. [De23, Ja07]. On the other hand, irreducible quotients/submodules of Jacquet modules could be a simpler object of study and is also useful for our applications. Our notion of derivatives is to study those irreucible quotients in certain form.

The main problem of this article is the following: given a collection of nice sequences of derivatives essentially square-integrable representations, can one find a canonical choice of an element among those that produce the same derivative. Here the term 'nice' refers to the sequences that arise from Bernstein-Zelevinsky derivatives. This article is to give an answer to this question. The results for those simple quotients may be regarded as generalizing the simpler cases of ladder representations in [LM14] and generic representations [Ch21].

In study of simple quotients of BZ derivatives is also related to a branching law problem of a sign representation for affine Hecke algebras of type A (see [Ch25, Appendix]). Some other applications such as on the Harish-Chandra modules for  $\operatorname{GL}_n(\mathbb{C})$  will be explored elsewhere, see e.g. [CW25].

Date: June 25, 2025.

1.1. Notations. Let  $G_n = \operatorname{GL}_n(F)$ , the general linear group over a non-Archimedean local field F. Fix a cuspidal representation  $\rho$ . We introduce basic notations:

Let ν : G<sub>n</sub> → C<sup>×</sup> be the character ν(g) = |det(g)|<sub>F</sub>, where |.|<sub>F</sub> is the norm for F.
For a, b ∈ Z with b − a ∈ Z<sub>>0</sub>, we call

$$(1.1) \qquad \qquad [a,b]_{\rho} := \left\{ \nu^a \rho, \dots, \nu^b \rho \right\}$$

to be a *segment*. For two segments  $\Delta$  and  $\Delta'$ , we write  $\Delta \cup \Delta'$  and  $\Delta \cap \Delta'$  for their set-theoretic union and intersection respectively.

We also set  $[a, a - 1]_{\rho} = \emptyset$  for  $a \in \mathbb{Z}$ . For a segment  $\Delta = [a, b]_{\rho}$ , we write  $a(\Delta) = \nu^{a}\rho$  and  $b(\Delta) = \nu^{b}\rho$ . We also write:

$$[a]_{\rho} := [a,a]_{\rho},$$

which is called a singleton segment. We may also write  $[\nu^a \rho, \nu^b \rho]$  for  $[a, b]_{\rho}$  and write  $[\nu^a \rho]$  for  $[a]_{\rho}$ . When we talk about long or short segments, we usually refer to the quantity b - a + 1 for a segment  $[a, b]_{\rho}$ .

- Let  $\operatorname{Seg}_{\rho}$  be the set of all segments. We also consider the empty set  $\emptyset$  to be in  $\operatorname{Seg}_{\rho}$ .
- A multisegment is a multiset of non-empty segments. Let  $\operatorname{Mult}_{\rho}$  be the set of all multisegments. We also consider the empty set  $\emptyset$  to be also in  $\operatorname{Mult}_{\rho}$ .
- Two segments Δ and Δ' are said to be *linked* if Δ ∪ Δ' is still a segment, and Δ ∉ Δ' and Δ' ∉ Δ. Otherwise, it is called to be not linked or unlinked.
- For  $\rho_1, \rho_2 \in \operatorname{Irr}^c$ , we write  $\rho_2 < \rho_1$  if  $\rho_1 \cong \nu^a \rho_2$  for some integer a > 0. For two segments  $\Delta_1, \Delta_2$ , we write  $\Delta_1 < \Delta_2$  if  $\Delta_1$  and  $\Delta_2$  are linked and  $b(\Delta_1) < b(\Delta_2)$ .
- For two multisegments m and n, we write m+n to be the union of two multisegments, counting multiplicities. For a multisegment m and a segment Δ, m + Δ = m + {Δ} if Δ is non-empty; and m + Δ = m if Δ is empty. The notions m n and m Δ are defined in a similar way.
- We say that a multisegment n is obtained from m by an elementary intersectionunion process if

$$\mathfrak{n} = \mathfrak{m} - \Delta - \Delta' + \Delta \cup \Delta' + \Delta \cap \Delta'$$

for a pair of linked segments  $\Delta$  and  $\Delta'$  in  $\mathfrak{m}$ .

- For two multisegments  $\mathfrak{m}_1$  and  $\mathfrak{m}_2$ , write  $\mathfrak{m}_2 \leq_Z \mathfrak{m}_1$  if  $\mathfrak{m}_2$  can be obtained by a sequence of elementary intersection-union operations from  $\mathfrak{m}_1$  (in the sense of [Ze80], see [Ch25]) or  $\mathfrak{m}_1 = \mathfrak{m}_2$ . In particular, if any pair of segments in  $\mathfrak{m}$  is unlinked, then  $\mathfrak{m}$  is a minimal element under  $\leq_Z$ . We shall equip Mult<sub> $\rho$ </sub> with the poset structure by  $\leq_Z$ .
- For a segment  $\Delta = [a, b]_{\rho}$ , define  $^{-}\Delta = [a + 1, b]_{\rho}$ . For a multisegment  $\mathfrak{m}$ , define  $^{-}\mathfrak{m} = \{^{-}\Delta : \Delta \in \mathfrak{m}, \Delta \text{ is not a singleton }\}$  (counting multiplicities).
- For each segment Δ, we shall denote by St(Δ) the corresponding essentially squareintegrable representation [Ze80].
- For any smooth representation  $\pi_1$  of  $G_{n_1}$  and smooth representation  $\pi_2$  of  $G_{n_2}$ , define  $\pi_1 \times \pi_2$  to be the normalized parabolic induction.
- Let  $\operatorname{Irr}_{\rho}(G_n)$  be the set of all irreducible representations of  $G_n$  which are irreducible quotients of  $\nu^{a_1}\rho \times \ldots \times \nu^{a_k}\rho$ , for some integers  $a_1, \ldots, a_k \in \mathbb{Z}$ . Let  $\operatorname{Irr}_{\rho} = \sqcup_n \operatorname{Irr}_{\rho}(G_n)$ .

1.2. Main results. Let  $N_i \subset G_n$  (depending on n) be the unipotent radical containing matrices of the form  $\begin{pmatrix} I_{n-i} & * \\ & I_i \end{pmatrix}$ . For a smooth representation  $\pi$  of  $G_n$ , we write  $\pi_{N_i}$  to be its Jacquet module.

For  $\pi \in \operatorname{Irr}_{\rho}(G_n)$  and a segment  $\Delta = [a, b]$ , there is at most one irreducible module  $\tau \in \operatorname{Irr}_{\rho}(G_{n-i})$  such that

$$\boxtimes \operatorname{St}(\Delta) \hookrightarrow \pi_{N_{b-a+1}}.$$

If such  $\tau$  exists, we denote such  $\tau$  by  $D_{\Delta}(\pi)$ . Otherwise, we set  $D_{\Delta}(\pi) = 0$ . We shall refer  $D_{\Delta}$  to be a *derivative*. Let  $\varepsilon_{\Delta}(\pi)$  be the largest integer k such that  $(D_{\Delta})^k(\pi) \neq 0$ .

A sequence of segments  $[a_1, b_1]_{\rho}, \ldots, [a_k, b_k]_{\rho}$  (where  $a_j, b_j \in \mathbb{Z}$ ) is said to be in an *ascending order* if for any  $i \leq j$ , either  $[a_i, b_i]_{\rho}$  and  $[a_j, b_j]_{\rho}$  are unlinked; or  $a_i < a_j$ . For a multisegment  $\mathfrak{n} \in \text{Mult}_{\rho}$ , which we write the segments in  $\mathfrak{n}$  in an ascending order  $\Delta_1, \ldots, \Delta_k$ . Define

$$D_{\mathfrak{n}}(\pi) := D_{\Delta_k} \circ \ldots \circ D_{\Delta_1}(\pi).$$

The derivative is independent of the ordering of an ascending sequence [Ch25]. In particular, one may choose the ordering such that  $a_1 \leq \ldots \leq a_k$ . We say that  $\mathfrak{n}$  is *admissible* to  $\pi$  if  $D_{\mathfrak{n}}(\pi) \neq 0$ . We refer the reader to [LM16, Ch22+b, Ch22+c] (and references therein) for more theory on derivatives.

For  $\pi \in \operatorname{Irr}_{\rho}$ , denote its *i*-th Bernstein-Zelevinsky derivative by  $\pi^{(i)}$  (see [Ze80, Ch25] for precise definitions and we shall not need this in this sequel). For a simple quotient  $\tau$  of  $\pi^{(i)}$ , define

$$\mathcal{S}(\pi,\tau) := \left\{ \mathfrak{n} \in \operatorname{Mult}_{\rho} : D_{\mathfrak{n}}(\pi) \cong \tau \right\}.$$

The ordering  $\leq_Z$  induces a partial ordering on  $\mathcal{S}(\pi,\tau)$ , and we shall regard  $\mathcal{S}(\pi,\tau)$  as a poset.

In [Ch25], we showed a combinatorial process, called *removal process*, in studying the effect of  $D_{\Delta}$ . Two applications of removal process are given below:

**Theorem 1.1.** (=Theorem 4.4) Let  $\pi \in \operatorname{Irr}_{\rho}$ . Let  $\tau$  be a simple quotient of  $\pi^{(i)}$  for some *i*. If  $\mathfrak{n}_1, \mathfrak{n}_2 \in \mathcal{S}(\pi, \tau)$  and  $\mathfrak{n}_1 \leq_Z \mathfrak{n}_2$ , then any  $\mathfrak{n}_3 \in \operatorname{Mult}_{\rho}$  satisfying  $\mathfrak{n}_1 \leq_Z \mathfrak{n}_3 \leq_Z \mathfrak{n}_2$  is also in  $\mathcal{S}(\pi, \tau)$ .

In other words,  $S(\pi, \tau)$  is *convex* in the sense of [St12, Section 3.1].

**Theorem 1.2.** (=Theorem 6.4) Let  $\pi \in \operatorname{Irr}_{\rho}$ . Let  $\tau$  be a simple quotient of  $\pi^{(i)}$  for some *i*. If  $S(\pi, \tau) \neq \emptyset$ , then  $S(\pi, \tau)$  has a unique minimal element with respect to  $\leq_Z$ .

The condition  $S(\pi, \tau) \neq \emptyset$  will be removed in [Ch22+b]. The non-emptyness condition will allow one to use Theorem 2.4 to transfer to combinatorial problems. As shown in [Ch21, Ch25], the socle of  $\pi^{(i)}$  is multiplicity-free, and the remaining problem to describe which representations appear in the socle of  $\pi^{(i)}$ . The minimal sequences give a canonical way to parametrize those representations, and in other words, we have a one-to-one correspondence between the admissible minimal sequences to  $\pi$  and the set of simple quotients of Bernstein-Zelevinsky derivatives of  $\pi$ .

For two segment case, there are some other criteria for a minimal sequence such as the nonoverlapping property and in terms of  $\eta_{\Delta}$  in Section 9. Such case is of particularly useful if one wants to 'split' some segments to carry out some inductive arguments and . We remark that there is no uniqueness for maximal elements in general. We give an example in Section 8.

1.3. Organization of this article. Section 2 recalls results on highest derivative multisegments and removal processes established in [Ch25]. Section 3 defines a notion of fine chains and fine chain orderings to facilitate comparisons with the Zelevinsky ordering. Section 4 shows the closedness property for  $S(\pi, \tau)$ . In Section 5, we shall introduce a notion of local minimizability, used to show the uniqueness of a minimal element in Section 6. Section 7 gives two examples of the unique minimal elements. Section 8 gives an example which

uniqueness of  $\leq_Z$ -maximality fails. Section 9 studies equivalent conditions for minimality in two segment cases.

1.4. Acknowledgements. This project is supported by the Research Grants Council of the Hong Kong Special Administrative Region, China (Project No: 17305223, 17308324) and NSFC grant for Excellent Young Scholar (Hong Kong and Macau) (Project No.: 12322120).

#### 2. Highest derivative multisegments and removal process

In this section, we recall some results in [Ch25].

2.1. More notations on multisegments. For an integer c, let  $\operatorname{Mult}_{\rho,c}^{a}$  be the subset of Mult<sub> $\rho$ </sub> containing all multisegments  $\mathfrak{m}$  such that any segment  $\Delta$  in  $\mathfrak{m}$  satisfies  $a(\Delta) \cong \nu^c \rho$ . The upperscript a in  $\operatorname{Mult}_{\rho,c}^{a}$  is for  $a(\Delta)$ .

For a multisegment  $\mathfrak{m}$  in Mult<sub> $\rho$ </sub> and an integer c, let  $\mathfrak{m}[c]$  be the submultisegment of  $\mathfrak{m}$ containing all the segments  $\Delta$  satisfying  $a(\Delta) \cong \nu^c \rho$ .

Fix an integer c. Let  $\Delta_1 = [c, b_1]_{\rho}, \Delta_2 = [c, b_2]_{\rho}$  be two non-empty segments. We write

 $\begin{array}{l} \Delta_1 \leq_c^a \Delta_2 \text{ if } b_1 \leq b_2, \text{ and write } \Delta_1 <_c^a \Delta_2 \text{ if } b_1 < b_2. \\ \text{For non-empty } \mathfrak{m}_1, \mathfrak{m}_2 \text{ in } \mathrm{Mult}_{\rho,c}^a, \text{ label the segments in } \mathfrak{m}_1 \text{ as: } \Delta_{1,k} \leq_c^a \ldots \leq_c^a \Delta_{1,2} \leq_c^a \\ \Delta_{1,1} \text{ and label the segments in } \mathfrak{m}_2 \text{ as: } \Delta_{2,r} \leq_c^a \ldots \leq_c^a \Delta_{2,2} \leq_c \Delta_{2,1}. \text{ We define the lexicographical ordering: } \mathfrak{m}_1 \leq_c^a \mathfrak{m}_2 \text{ if } k \leq r \text{ and, for any } i \leq k, \Delta_{1,i} \leq_c^a \Delta_{2,i}. \end{array}$  $\mathfrak{m}_1 <^a_c \mathfrak{m}_2$  if  $\mathfrak{m}_1 \leq^a_c \mathfrak{m}_2$  and  $\mathfrak{m}_1 \neq \mathfrak{m}_2$ .

2.2. Highest derivative multisegments. A multisegment  $\mathfrak{m}$  is said to be at the point  $\nu^c \rho$  if any segment  $\Delta$  in  $\mathfrak{m}$  takes the form  $[c,b]_\rho$  for some  $b \geq c$ . For  $\pi \in \operatorname{Irr}_{\rho}$ , define  $\mathfrak{mppt}^{a}(\pi, c)$  to be the maximal multisegment in  $\mathrm{Mult}^{a}_{\rho,c}$  such that  $D_{\mathfrak{mppt}^{a}(\pi,c)}(\pi) \neq 0$ . The maximality is determined by the ordering  $\leq_c^a$  defined above. Define the highest derivative multisegment of  $\pi \in \operatorname{Irr}_{\rho}$  to be

$$\mathfrak{hd}(\pi) := \sum_{c \in \mathbb{Z}} \mathfrak{mpt}^a(\pi, c).$$

It is shown in [Ch25] that  $D_{\mathfrak{ho}(\pi)}(\pi)$  is the highest derivative of  $\pi$  in the sense of [Ze80].

2.3. Removal process. We write  $[a,b]_{\rho} \prec^{L} [a',b']_{\rho}$  if either a < a'; or a = a' and b < b'. Here L means to compare on the 'left' value a and we avoid to further use a for confusing with previous notations. A non-empty segment  $\Delta = [a, b]_{\rho}$  is said to be *admissible* to a multisegment  $\mathfrak{h}$  if there exists a segment of the form  $[a, c]_{\rho}$  in  $\mathfrak{h}$  for some  $c \geq b$ . We now recall the removal process.

An *infinity multisegment*, denoted by  $\infty$ , is just a symbol which will be used to indicate the situation of non-admissibility.

**Definition 2.1.** [Ch25] Let  $\mathfrak{h} \in \text{Mult}_{\rho}$  and let  $\Delta = [a, b]_{\rho}$  be admissible to  $\mathfrak{h}$ . The removal process on  $\mathfrak{h}$  by  $\Delta$  is an algorithm to carry out the following steps:

- (1) Pick a shortest segment  $[a, c]_{\rho}$  in  $\mathfrak{h}[a]$  satisfying  $b \leq c$ . Set  $\Delta_1 = [a, c]_{\rho}$ . Set  $a_1 = a$ and  $b_1 = c$ .
- (2) One recursively finds the  $\prec^L$ -minimal segment  $\Delta_i = [a_i, b_i]_{\rho}$  in  $\mathfrak{h}$  such that  $a_{i-1} < a_i$ and  $b_i < b_{i-1}$ . The process stops if one can no longer find those segments.
- (3) Let  $\Delta_1, \ldots, \Delta_r$  be all those segments. For  $1 \leq i < r$ , define  $\Delta_i^{tr} = [a_{i+1}, b_i]_{\rho}$  and  $\Delta_r^{tr} = [b+1, b_r]_{\rho}$  (possibly empty).

(4) Define

$$\mathfrak{r}(\Delta, \mathfrak{h}) := \mathfrak{h} - \sum_{i=1}^{r} \Delta_i + \sum_{i=1}^{r} \Delta_i^{tr}.$$

We call  $\Delta_1, \ldots, \Delta_r$  to be the removal sequence for  $(\Delta, \mathfrak{h})$ . We also define  $\Upsilon(\Delta, \mathfrak{h}) = \Delta_1$ , the first segment in the removal sequence. If  $\Delta$  is not admissible to  $\mathfrak{h}$ , we set  $\mathfrak{r}(\Delta, \mathfrak{h}) = \infty$ , the infinity multisegment. We also set  $\mathfrak{r}(\Delta, \infty) = \infty$ .

**Remark 2.2.** A multisegment  $\mathfrak{h}$  is called *generic* if any two segments in  $\mathfrak{h}$  are unlinked. The special feature in this case is that  $\operatorname{St}(\mathfrak{h})$  is generic and  $\mathfrak{hd}(\operatorname{St}(\mathfrak{h})) = \mathfrak{h}$ . In such case, for  $\Delta \in \operatorname{Seg}_{\rho}$ , it is shown in [Ch21] that  $D_{\Delta}(\pi)$  is generic. On the other hand,  $\mathfrak{r}(\Delta, \mathfrak{h})$  coincides with the generic multisegment which has the same cuspidal support as  $D_{\Delta}(\pi)$ .

2.4. **Computations on removal process.** We recall the following properties for computations:

**Lemma 2.3.** [Ch25] Let  $\mathfrak{h} \in \text{Mult}_{\rho}$  and let  $\Delta, \Delta' \in \text{Seg}_{\rho}$  be admissible to  $\mathfrak{h}$ . Then the followings hold:

- (1) Let  $\mathfrak{h}^* = \mathfrak{h} \Upsilon(\Delta, \mathfrak{h}) + {}^{-}\Upsilon(\Delta, \mathfrak{h})$ . Then  $\mathfrak{r}(\Delta, \mathfrak{h}) = \mathfrak{r}({}^{-}\Delta, \mathfrak{h}^*)$ .
- (2) Write  $\Delta = [a, b]_{\rho}$ . For any a' < a,  $\mathfrak{r}(\Delta, \mathfrak{h})[a'] = \mathfrak{h}[a']$ .
- (3) If  $\Delta \in \mathfrak{h}$ , then  $\mathfrak{r}(\Delta, \mathfrak{h}) = \mathfrak{h} \Delta$ .
- (4) Suppose  $a(\Delta) = a(\Delta')$ . Then

 $\Upsilon(\Delta, \mathfrak{h}) + \Upsilon(\Delta', \mathfrak{r}(\Delta, \mathfrak{h})) = \Upsilon(\Delta', \mathfrak{h}) + \Upsilon(\Delta, \mathfrak{r}(\Delta', \mathfrak{h})).$ 

(5) If  $\Delta, \Delta'$  are unlinked, then  $\mathfrak{r}(\Delta', \mathfrak{r}(\Delta, \mathfrak{h})) = \mathfrak{r}(\Delta, \mathfrak{r}(\Delta', \mathfrak{h})).$ 

2.5. Removal process for multisegments. For  $\mathfrak{h} \in \text{Mult}_{\rho}$ , and a multisegment  $\mathfrak{m} = \{\Delta_1, \ldots, \Delta_r\} \in \text{Mult}_{\rho}$  written in an ascending order (defined in Section 1.2), define:

$$\mathfrak{r}(\mathfrak{m},\mathfrak{h}) = \mathfrak{r}(\Delta_r,\ldots,\mathfrak{r}(\Delta_1,\mathfrak{h})\ldots).$$

We say that  $\mathfrak{m}$  is *admissible* to  $\mathfrak{h}$  if  $\mathfrak{r}(\mathfrak{m}, \mathfrak{h}) \neq \infty$ .

For  $\pi \in \operatorname{Irr}_{\rho}$  and  $\mathfrak{m} \in \operatorname{Mult}_{\rho}$ , define  $\mathfrak{r}(\mathfrak{m}, \pi) := \mathfrak{r}(\mathfrak{m}, \mathfrak{hd}(\pi))$ . The relation to derivatives is the following:

**Theorem 2.4.** [Ch25] Let  $\pi \in \operatorname{Irr}_{\rho}$ . Let  $\mathfrak{m}, \mathfrak{m}' \in \operatorname{Mult}_{\rho}$  be admissible to  $\pi$ . Then  $\mathfrak{m}, \mathfrak{m}'$  are admissible to  $\mathfrak{ho}(\pi)$ , and furthermore,  $D_{\mathfrak{m}}(\pi) \cong D_{\mathfrak{m}'}(\pi)$  if and only if  $\mathfrak{r}(\mathfrak{m}, \pi) = \mathfrak{r}(\mathfrak{m}', \pi)$ .

2.6. More relations to derivatives. For  $\mathfrak{h} \in \mathrm{Mult}_{\rho}$  and  $\Delta = [a, b]_{\rho} \in \mathrm{Seg}_{\rho}$ , set

$$\varepsilon_{\Delta}(\mathfrak{h}) = |\left\{\widetilde{\Delta} \in \mathfrak{h}[a] : \Delta \subset \widetilde{\Delta}\right\}|.$$

**Theorem 2.5.** [Ch25, Theorem 6.20] Let  $\pi \in \operatorname{Irr}_{\rho}$ . Let  $\Delta \in \operatorname{Seg}_{\rho}$  be admissible to  $\pi$ . Let  $\Delta' \in \operatorname{Seg}_{\rho}$ . Suppose either  $a(\Delta') > a(\Delta)$ ; or  $\Delta'$  and  $\Delta$  are unlinked. Then  $\varepsilon_{\Delta'}(D_{\Delta}(\pi)) = \varepsilon_{\Delta'}(\mathfrak{r}(\Delta, \pi))$ .

#### 3. FINE CHAINS

Recall that  $S(\pi, \tau)$  is defined in Section 1.2. We introduce a notion of fine chains in Definition 3.5 in order to give an effective comparison of the effect of two removal processes (Lemma 3.9).

3.1. Basic idea on the proof for Theorem 4.4. Let  $\mathfrak{h} \in \operatorname{Mul}_{\rho}$ . We consider two segments in the form  $\Delta = [a, b]_{\rho}$  and  $\Delta' = [a + 1, b']_{\rho}$  for b < b'. In such case, let  $\widetilde{\Delta} = \Delta \cup \Delta' = [a, b']_{\rho}$  and  $\widetilde{\Delta}' = \Delta \cap \Delta' = [a + 1, b]_{\rho}$ . Let  $\Omega = \Upsilon(\Delta, \mathfrak{h})$  and let  $\widetilde{\Omega} = \Upsilon(\widetilde{\Delta}, \mathfrak{h})$ . Let

$$\mathfrak{h}^* = \mathfrak{h} - \Omega + {}^{-}\Omega, \quad \widetilde{\mathfrak{h}} = \mathfrak{h} - \widetilde{\Omega} + {}^{-}\widetilde{\Omega}.$$

Applying Lemma 2.3, we have

$$\mathfrak{r}(\{\Delta',\Delta\}\,,\mathfrak{h})=\mathfrak{r}(\{\Delta',{}^{-}\Delta\}\,,\mathfrak{h}^*)$$

and

$$\mathfrak{r}(\left\{\widetilde{\Delta}',\widetilde{\Delta}\right\},\mathfrak{h}) = \mathfrak{r}(\left\{\widetilde{\Delta}',^{-}\widetilde{\Delta}\right\},\widetilde{\mathfrak{h}}).$$

Note that  $\{\Delta', {}^{-}\Delta\} = \{\widetilde{\Delta}', {}^{-}\widetilde{\Delta}\}$ . Then the following statements are equivalent:

- $\begin{array}{ll} (1) \ \mathfrak{r}(\{\Delta,\Delta'\}\,,\mathfrak{h})=\mathfrak{r}(\{\widetilde{\Delta},\widetilde{\Delta}'\},\mathfrak{h});\\ (2) \ \mathfrak{h}^*=\widecheck{\mathfrak{h}}; \end{array}$
- (3)  $\Omega = \widetilde{\Omega}$ .

The general case for the effect of intersection-union processes needs some modifications for the above consideration and we shall focus on the condition (3). In particular, Sections 3.2 and 3.3 will improve Lemma 2.3(1) to do some multiple 'cutting-off' on starting points.

3.2. Multiple removal of starting points. Let  $\mathfrak{h} \in \text{Mult}_{\rho}$ . Let  $\mathfrak{n} \in \text{Mult}_{\rho}$ . Let a be the smallest integer such that  $\mathfrak{n}[a] \neq \emptyset$ . Write  $\mathfrak{n}[a] = \{\Delta_1, \ldots, \Delta_k\}$ .

(1) Suppose  $\mathfrak{n}[a]$  is admissible to  $\mathfrak{h}$ . Let  $\mathfrak{r}_i = \mathfrak{r}(\{\Delta_i, \ldots, \Delta_1\}, \mathfrak{h})$  and  $\mathfrak{r}_0 = \mathfrak{h}$ . Define

 $\mathfrak{fs}(\mathfrak{n},\mathfrak{h}) = \{\Upsilon(\Delta_1,\mathfrak{r}_0),\ldots,\Upsilon(\Delta_k,\mathfrak{r}_{k-1})\}.$ 

(fs refers to first segments.)

(2) Suppose  $\mathfrak{n}[a]$  is not admissible to  $\mathfrak{h}$ . Define  $\mathfrak{fs}(\mathfrak{n},\mathfrak{h}) = \emptyset$ .

**Lemma 3.1.** The above definition  $\mathfrak{s}(\mathfrak{n},\mathfrak{h})$  is well-defined i.e. independent of an ordering for the segments in  $\mathfrak{n}[a]$ .

*Proof.* One switches a consecutive pair of segments each time, and then applies Lemmas 2.3(4) and 2.3(5).

Following from definitions, we also have:

**Lemma 3.2.** With the notations as above,

$$\mathfrak{s}(\mathfrak{n},\mathfrak{h}) = \mathfrak{s}(\mathfrak{n}[a],\mathfrak{h}) = \mathfrak{s}(\mathfrak{n}[a],\mathfrak{h}[a]).$$

We define a truncation of  $\mathfrak{h}$ :

(3.2)  $\operatorname{trr}(\mathfrak{n},\mathfrak{h}) = \mathfrak{h} - \mathfrak{fs}(\mathfrak{n},\mathfrak{h}) + {}^{-}(\mathfrak{fs}(\mathfrak{n},\mathfrak{h})),$ 

and a truncation of  $\mathfrak{n}$ :

$$\mathfrak{trd}(\mathfrak{n},\mathfrak{h}) = \mathfrak{n} - \mathfrak{n}[a] + {}^{-}(\mathfrak{n}[a])$$

Here we use r in  $\mathfrak{trr}$  for the derivative 'resultant' multisegment and d for  $\mathfrak{trd}$  for 'taking the derivative for the multisegment  $\mathfrak{n}$ '.

**Example 3.3.** Let  $\mathfrak{h} = \{[0,3]_{\rho}, [1,2]_{\rho}, [1,4]_{\rho}, [1,5]_{\rho}, [2,3]_{\rho}\}$ . Let  $\mathfrak{n} = \{[1,3]_{\rho}, [1,5]_{\rho}, [2]_{\rho}\}$ .

 $\mathbf{6}$ 



The two red bullets in  $\mathfrak{h}$  are 'truncated' to obtain  $\mathfrak{trr}(\mathfrak{n},\mathfrak{h}) = \{[0,3]_{\rho}, [1,2]_{\rho}, [2,4]_{\rho}, [2,5]_{\rho}, [2,3]_{\rho}\}$ . We also have  $\mathfrak{fs}(\mathfrak{n},\mathfrak{h}) = \{[1,4]_{\rho}, [1,5]_{\rho}\}$  and  $\mathfrak{trd}(\mathfrak{n},\mathfrak{h}) = \{[2,3]_{\rho}, [2,5]_{\rho}, [2]_{\rho}\}$ .

**Lemma 3.4.** (multiple removal of starting points, c.f. Lemma 2.3(1)) Let  $\mathfrak{n}, \mathfrak{h}, \mathfrak{a}$  be as above. Then

$$\mathfrak{r}(\mathfrak{n},\mathfrak{h}) = \mathfrak{r}(\mathfrak{trd}(\mathfrak{n},\mathfrak{h}),\mathfrak{trr}(\mathfrak{n},\mathfrak{h}))$$

*Proof.* Write  $\mathfrak{n}[a] = \{\overline{\Delta}_1, \dots, \overline{\Delta}_k\}$ . Relabeling if necessary,  $\overline{\Delta}_1$  is the shortest segment in  $\mathfrak{n}[a]$ . Let

$$\mathfrak{h}_1^* = \mathfrak{h} - \left\{\overline{\Delta}_1\right\} + \left\{\overline{\Delta}_1\right\}$$

We observe that:

$$\begin{aligned} \mathfrak{r}(\mathfrak{n}[a],\mathfrak{h}) &= \mathfrak{r}(\left\{\overline{\Delta}_2,\ldots,\overline{\Delta}_k\right\},\mathfrak{r}(\overline{\Delta}_1,\mathfrak{h})) \\ &= \mathfrak{r}(\left\{\overline{\Delta}_2,\ldots,\overline{\Delta}_k\right\},\mathfrak{r}(^{-}\overline{\Delta}_1,\mathfrak{h}_1^*)) \\ &= \mathfrak{r}(^{-}\overline{\Delta}_1,\mathfrak{r}(\left\{\overline{\Delta}_2,\ldots,\overline{\Delta}_k\right\},\mathfrak{h}_1^*)) \\ &= \mathfrak{r}(^{-}\overline{\Delta}_1,\mathfrak{r}(\left\{^{-}\overline{\Delta}_2,\ldots,^{-}\overline{\Delta}_k\right\},\mathfrak{trr}(\mathfrak{n},\mathfrak{h}))) \\ &= \mathfrak{r}(^{-}(\mathfrak{n}[a]),\mathfrak{trr}(\mathfrak{n},\mathfrak{h})), \end{aligned}$$

where the second equation follows from Lemma 2.3(1), the first, third and last equations follow from Lemma 2.3(5), and the forth equation follows from the induction hypothesis (where the basic case is again Lemma 2.3(1)).

The lemma then follows by applying  $\mathfrak{r}(\mathfrak{n} - \mathfrak{n}[a], .)$  on the first and last terms.

### 3.3. Fine chains.

**Definition 3.5.** For  $c \in \mathbb{Z}$ , we modify the ordering  $\langle a \\ c \\ c \\ on Mult^a_{\rho,c} \cup \{\infty\}$  as follows. For  $\mathfrak{p}_1, \mathfrak{p}_2$  in  $Mult^a_{\rho,c} \cup \{\infty\}$ , if  $\mathfrak{p}_1 \neq \infty$  and  $\mathfrak{p}_2 = \infty$ , we also write  $\mathfrak{p}_1 \langle a \\ c \\ \mathfrak{p}_2$ . If  $\mathfrak{p}_1 = \mathfrak{p}_2 = \infty$ , we write  $\mathfrak{p}_1 \langle a \\ c \\ \mathfrak{p}_2$ .

**Definition 3.6.** (Collections of first segments in the removal sequence) Let  $\mathfrak{h} \in \text{Mult}_{\rho}$ . Let  $\mathfrak{n} \in \text{Mult}_{\rho}$ . Set  $\mathfrak{n}_0 = \mathfrak{n}$  and  $\mathfrak{h}_0 = \mathfrak{h}$ . We recursively define:

$$\mathfrak{h}_i = \mathfrak{trr}(\mathfrak{h}_{i-1}, \mathfrak{n}_{i-1}), \quad \mathfrak{n}_i = \mathfrak{trd}(\mathfrak{h}_{i-1}, \mathfrak{n}_{i-1}).$$

The sequence of multisegments

$$\mathfrak{fs}(\mathfrak{n}_0,\mathfrak{h}_0),\mathfrak{fs}(\mathfrak{n}_1,\mathfrak{h}_1),\ldots$$

 $\overline{7}$ 

is called the *fine chain* for  $(\mathfrak{n}, \mathfrak{h})$ . Since we usually fix  $\mathfrak{h}$  and vary  $\mathfrak{n}$  in our use of fine chains, we shall denote the fine chain by  $fc_{\mathfrak{h}}(\mathfrak{n})$ . It follows from the definition that  $\mathfrak{fs}(\mathfrak{n}_i, \mathfrak{h}_i), \mathfrak{fs}(\mathfrak{n}_{i+1}, \mathfrak{h}_{i+1}), \ldots$  is also the fine chain for  $(\mathfrak{n}_i, \mathfrak{h}_i)$ .

**Example 3.7.** Let  $\mathfrak{h} = \{[0,4]_{\rho}, [1,5]_{\rho}\}.$ 

• Let  $\mathfrak{n} = \{[0,1]_{\rho}, [1,2]_{\rho}\}$ . Then the fine chain for  $(\mathfrak{n},\mathfrak{h})$  takes the form:

 $\{[0,4]_{\rho}\}, \{[1,4]_{\rho}, [1,5]_{\rho}\}, \{[2,4]_{\rho}\}.$ 

Let n = {[0,2]<sub>ρ</sub>, [1]<sub>ρ</sub>}. Then the fine chain for (n, h) is the same as the previous one.

**Definition 3.8.** Let  $\mathfrak{h} \in \operatorname{Mult}_{\rho}$ . We say that two fine chains  $\operatorname{fc}_{\mathfrak{h}}(\mathfrak{n})$  and  $\operatorname{fc}_{\mathfrak{h}}(\mathfrak{n}')$  coincide if

- (1)  $\mathfrak{r}(\mathfrak{n},\mathfrak{h}), \mathfrak{r}(\mathfrak{n}',\mathfrak{h}) \neq \infty$ ; and
- (2) the two sequences  $fc_{\mathfrak{h}}(\mathfrak{n})$  and  $fc_{\mathfrak{h}}(\mathfrak{n}')$  are equal.

**Lemma 3.9.** Let  $\mathfrak{h} \in \text{Mult}_{\rho}$ . Let  $\mathfrak{n}, \mathfrak{n}' \in \text{Mult}_{\rho}$ . Then  $\mathfrak{r}(\mathfrak{n}, \mathfrak{h}) = \mathfrak{r}(\mathfrak{n}', \mathfrak{h}) \neq \infty$  if and only if the fine chains  $\text{fc}_{\mathfrak{h}}(\mathfrak{n})$  and  $\text{fc}_{\mathfrak{h}}(\mathfrak{n}')$  coincide.

*Proof.* We write fine chains  $fc_{\mathfrak{h}}(\mathfrak{n})$  and  $fc_{\mathfrak{h}}(\mathfrak{n}')$  with notations in Definition 3.6 as:

 $\mathfrak{fs}(\mathfrak{n}_0,\mathfrak{h}_0),\mathfrak{fs}(\mathfrak{n}_1,\mathfrak{h}_1),\ldots$ 

and

 $\mathfrak{fs}(\mathfrak{n}_0',\mathfrak{h}_0'),\mathfrak{fs}(\mathfrak{n}_1',\mathfrak{h}_1'),\ldots$ 

with  $\mathfrak{n}_0 = \mathfrak{n}, \, \mathfrak{n}'_0 = \mathfrak{n}', \, \mathfrak{h}_0 = \mathfrak{h}'_0 = \mathfrak{h}.$ 

For the only if direction, Lemma 2.3(2) implies that  $\mathfrak{h}_i = \mathfrak{h}'_i$  for all *i*. Then, from the construction of  $\mathfrak{h}_i$  and  $\mathfrak{h}'_i$ , we have that  $\mathfrak{fs}(\mathfrak{n}_{i-1},\mathfrak{h}_{i-1}) = \mathfrak{fs}(\mathfrak{n}'_{i-1},\mathfrak{h}'_{i-1})$ . In other words, the fine chains for  $(\mathfrak{n},\mathfrak{h})$  and  $(\mathfrak{n}',\mathfrak{h})$  coincide.

For the if direction, since the fine chains coincide, we must have  $\mathfrak{h}_i = \mathfrak{h}'_i$  by (3.2). In particular,  $\mathfrak{r}(\mathfrak{n},\mathfrak{h}) = \mathfrak{r}(\mathfrak{n}',\mathfrak{h})$  as desired.

3.4. Fine chain ordering. Multisegments  $\mathfrak{n}$  and  $\mathfrak{n}'$  are said to be of the same cuspidal support if  $\bigcup_{\Delta \in \mathfrak{n}} \Delta = \bigcup_{\Delta \in \mathfrak{n}'} \Delta$  (counting multiplicities).

**Definition 3.10.** Let  $\mathfrak{n}, \mathfrak{n}' \in \operatorname{Mult}_{\rho}$  be of the same cuspidal support. Let  $\mathfrak{h} \in \operatorname{Mult}_{\rho}$ . Suppse  $\mathfrak{r}(\mathfrak{n}, \mathfrak{h}) \neq \infty$  and  $\mathfrak{r}(\mathfrak{n}', \mathfrak{h}) \neq \infty$ . Write  $fc_{\mathfrak{h}}(\mathfrak{n})$  as  $\mathfrak{s}_1, \mathfrak{s}_2, \ldots$  and write  $fc_{\mathfrak{h}}(\mathfrak{n}')$  as  $\mathfrak{s}'_1, \mathfrak{s}'_2, \ldots$  Similar to the notations in Definition 3.6, set inductively  $\mathfrak{n}_i = \mathfrak{n}_{i-1} - \mathfrak{s}_i + -\mathfrak{s}_i$  and  $\mathfrak{n}'_i = \mathfrak{n}'_{i-1} - \mathfrak{s}'_i + -\mathfrak{s}'_i$ , where  $\mathfrak{n}_0 = \mathfrak{n}$  and  $\mathfrak{n}'_0 = \mathfrak{n}'$ . Let  $c_i$  (resp.  $c'_i$ ) be the smallest integer such that  $\mathfrak{n}_i[c_i] \neq \emptyset$  (resp.  $\mathfrak{n}'_i[c'_i] \neq \emptyset$ ).

We define  $\mathfrak{n} <^{fc} \mathfrak{n}'$ , called the *fine chain ordering*, if there exists some *i* such that for any  $j < i, \mathfrak{s}_j = \mathfrak{s}'_j$  and

 $\mathfrak{s}_i <^a_{c_{i-1}} \mathfrak{s}'_i.$ 

We write  $\mathfrak{n} \leq f^c \mathfrak{n}'$  if either  $\mathfrak{n} < f^c \mathfrak{n}'$  or  $fc_{\mathfrak{h}}(\mathfrak{n}) = fc_{\mathfrak{h}}(\mathfrak{n}')$ . Note that  $\leq f^c$  is transitive.

### 4. CLOSURE UNDER INTERSECTION-UNION PROCESS

The main result in this section is Theorem 4.4, which gives a combinatorial structure of  $S(\pi, \tau)$ . Theorem 2.4 transfers the problem on the convex structure of  $S(\pi, \tau)$  to study explicit combinatorics on the removal process.

#### 4.1. Effect from intersection-union process.

**Lemma 4.1.** Let  $\mathfrak{h} \in \operatorname{Mult}_{\rho}$ . Let  $\mathfrak{m}_1$  be in  $\operatorname{Mult}^a_{\rho,c}$ . Let  $\mathfrak{m}_2 \in \operatorname{Mult}^a_{\rho,c}$  be obtained from  $\mathfrak{m}_1$  by replacing one segment in  $\mathfrak{m}_1$  with a longer segment of the form  $[c, b]_{\rho}$  for some  $b \in \mathbb{Z}$ . Then

$$\mathfrak{fs}(\mathfrak{m}_1,\mathfrak{h})\leq^a_c\mathfrak{fs}(\mathfrak{m}_2,\mathfrak{h}).$$

*Proof.* The only difference between  $\mathfrak{m}_1$  and  $\mathfrak{m}_2$  is on one segment. We can arrange those segments to be the last ones in the process of obtaining  $\mathfrak{fs}(\mathfrak{m}_1,\mathfrak{h})$  and  $\mathfrak{fs}(\mathfrak{m}_2,\mathfrak{h})$  respectively by Lemma 3.1. Thus the only difference between  $\mathfrak{fs}(\mathfrak{m}_1,\mathfrak{h})$  and  $\mathfrak{fs}(\mathfrak{m}_2,\mathfrak{h})$  is only one segment. The remaining one follows from the definition of  $\Upsilon$  (for picking the last segments in  $\mathfrak{fs}(\mathfrak{m}_1,\mathfrak{h})$  and  $\mathfrak{fs}(\mathfrak{m}_2,\mathfrak{h})$ ) and  $\mathfrak{fs}(\mathfrak{m}_2,\mathfrak{h})$ .

**Lemma 4.2.** Let  $\mathfrak{h} \in \text{Mult}_{\rho}$ . Fix  $\mathfrak{n} \in \text{Mult}_{\rho}$ . Let  $\mathcal{N} = \mathcal{N}(\mathfrak{n})$  be the set of all multisegments of the same cuspidal support as  $\mathfrak{n}$ . Then, for  $\mathfrak{n}', \mathfrak{n}'' \in \mathcal{N}$ ,

$$\mathfrak{n}' \leq_Z \mathfrak{n}'' \implies \mathfrak{n}'' \leq^{fc} \mathfrak{n}'.$$

*Proof.* By the transitivity of  $\leq_Z$ , we reduce to the case that  $\mathfrak{n}'$  is obtained from  $\mathfrak{n}''$  by an elementary intersection-union operation. Let  $\Delta_1$  and  $\Delta_2$  be the two linked segments involved in the elementary intersection-union operation. Relabeling if necessary, we write:

$$\Delta_1 = [a_1, b_1]_{\rho}, \quad \Delta_2 = [a_2, b_2]_{\rho},$$

with  $a_1 < a_2$  and  $b_1 < b_2$ .

We again write  $f_{c_b}(\mathfrak{n}')$  as  $\mathfrak{s}'_1, \mathfrak{s}'_2, \ldots$  and write  $f_{c_b}(\mathfrak{n}'')$  as  $\mathfrak{s}''_1, \mathfrak{s}''_2, \ldots$ . Similar to notations in Definition 3.6, set  $\mathfrak{n}'_i = \mathfrak{n}'_{i-1} - \mathfrak{s}'_i + -\mathfrak{s}'_i$  and  $\mathfrak{n}''_i = \mathfrak{n}''_{i-1} - \mathfrak{s}''_i + -\mathfrak{s}''_i$ . Again let  $c_i$  be the smallest integer such the  $\mathfrak{n}'_i[c_i] \neq \emptyset$ . It is straighforward to see from the intersection-union operation that  $\mathfrak{n}''_i[c_i]$  is obtained from  $\mathfrak{n}'_i[c_i]$  by replacing a segment with a longer one (of the form  $[c_i, b]_{\rho}$ ). Thus now Lemma 4.1 implies that  $\mathfrak{n}'' \leq f^c \mathfrak{n}'$ .

**Theorem 4.3.** Let  $\mathfrak{n}, \mathfrak{n}' \in \text{Mult}_{\rho}$ . Suppose  $\mathfrak{n}' \leq_Z \mathfrak{n}$ . Let  $\mathfrak{n}'' \in \text{Mult}_{\rho}$  such that

$$\mathfrak{n}' \leq_Z \mathfrak{n}'' \leq_Z \mathfrak{n}.$$

Then, if  $\mathfrak{r}(\mathfrak{n},\mathfrak{h}) = \mathfrak{r}(\mathfrak{n}',\mathfrak{h})$ , then  $\mathfrak{r}(\mathfrak{n},\mathfrak{h}) = \mathfrak{r}(\mathfrak{n}'',\mathfrak{h})$ .

*Proof.* If  $\mathfrak{r}(\mathfrak{n},\mathfrak{h}) \neq \mathfrak{r}(\mathfrak{n}'',\mathfrak{h})$ , Lemmas 3.9 and 4.2 imply that  $\mathfrak{n}'' <^{fc} \mathfrak{n}$ . By Lemma 4.2 again,  $\mathfrak{n}' <_Z \mathfrak{n}$ . Now the transitivity of  $<^{fc}$  implies that  $\mathfrak{n}' <^{fc} \mathfrak{n}$ . However, Lemma 3.9 then implies  $\mathfrak{r}(\mathfrak{n}',\mathfrak{h}) \neq \mathfrak{r}(\mathfrak{n},\mathfrak{h})$ , giving a contradiction.

We translate the combinatorial statement in Theorem 4.3 to its representation-theoretic counterpart:

**Theorem 4.4.** Let  $\pi \in \operatorname{Irr}_{\rho}$  and let  $\tau$  be a simple quotient of  $\pi^{(i)}$ . Recall that  $S(\pi, \tau)$  is defined in Section 1.2. Let  $\mathfrak{n}, \mathfrak{n}' \in S(\pi, \tau)$  with  $\mathfrak{n}' \leq_Z \mathfrak{n}$ . For any  $\mathfrak{n}'' \in \operatorname{Mult}_{\rho}$  such that  $\mathfrak{n}' \leq_Z \mathfrak{n}'' \leq_Z \mathfrak{n}$ , we have  $\mathfrak{n}'' \in S(\pi, \tau)$ .

*Proof.* This follows from Theorem 2.4 and Theorem 4.3.

We also have the following combinatorial consequence:

Corollary 4.5. We use the notations in Lemma 4.2. Let

$$\widetilde{\mathcal{N}}:=\left\{\mathfrak{n}\in\mathcal{N}:\mathfrak{r}(\mathfrak{n},\mathfrak{h})
eq\infty
ight\}.$$

We define an equivalence relation  $\sim on \widetilde{\mathcal{N}}$  by:  $\mathfrak{n} \sim \mathfrak{n}'$  if and only if  $\mathfrak{r}(\mathfrak{n},\mathfrak{h}) = \mathfrak{r}(\mathfrak{n}',\mathfrak{h})$ . Define  $\preceq_Z on \widetilde{\mathcal{N}} / \sim by$ : for  $N, N' \in \widetilde{\mathcal{N}} / \sim$ , write  $N \preceq_Z N'$  if there exists  $\mathfrak{n} \in N$  and  $\mathfrak{n}' \in N'$  such that  $\mathfrak{n} \leq_Z \mathfrak{n}'$ . We similarly define the notion  $\preceq^{f_c}$  on  $\widetilde{\mathcal{N}}$  by replacing  $\leq_Z$  with  $\leq^{f_c}$ . Then, the following holds:

- Both  $\leq_Z$  and  $\leq^{fc}$  define a well-defined poset structure on  $\widetilde{\mathcal{N}}/\sim$ .

*Proof.* For the first bullet, the only non-evident part is the antisymmetry, which indeed follows from Lemmas 3.9 and 4.2. The second bullet is a direct consequence on Lemma 4.2.  $\Box$ 

#### 5. MINIMIZABILITY

### 5.1. Basic example on minimality.

**Example 5.1.** Let  $\mathfrak{h} = \{[0,5]_{\rho}, [3,8]_{\rho}\}$ . Let  $\mathfrak{n} = \{[0,3]_{\rho}, [3,4]_{\rho}\}$ . Then  $\mathfrak{r}([0,3]_{\rho},\mathfrak{h}) = \{[4,5]_{\rho}, [3,8]_{\rho}\}$  and so  $\mathfrak{r}([3,4]_{\rho}, \mathfrak{r}([0,3]_{\rho},\mathfrak{h})) = \{[4,8]_{\rho}, [5]_{\rho}\}$ . Note that the segmet  $[5]_{\rho}$  coming from truncating the segment  $[4,5]_{\rho}$  in  $\mathfrak{r}([0,3]_{\rho},\mathfrak{h})$  and the segment  $[4,5]_{\rho}$  indeed comes from truncating the segment  $[0,5]_{\rho}$ . One wonders if one can 'combine' these two effects. Indeed, if one could consider  $\mathfrak{n}' = \{[0,4]_{\rho}, [3]_{\rho}\}$ , then  $\mathfrak{r}([3]_{\rho},\mathfrak{h}) = \{[0,5]_{\rho}, [4,8]_{\rho}\}$  and  $\mathfrak{r}([0,4]_{\rho},\mathfrak{r}([3]_{\rho},\mathfrak{h})) = \{[5]_{\rho}, [4,8]_{\rho}\}$ . In the last removal process,  $[5]_{\rho}$  is obtained directly from truncating  $[0,5]_{\rho}$  once.

For convenience, we define a multisegment analogue of  $\mathcal{S}(\pi, \tau)$ . For  $\mathfrak{h}, \mathfrak{p} \in \text{Mult}_{\rho}$ ,

$$\mathcal{S}'(\mathfrak{h},\mathfrak{p}) = \{\mathfrak{m} \in \mathrm{Mult}_{\rho} : \mathfrak{r}(\mathfrak{m},\mathfrak{h}) = \mathfrak{p}\}.$$

The above example shows that  $\mathfrak{n}$  is not  $\leq_Z$ -minimal in  $\mathcal{S}'(\mathfrak{h}, \mathfrak{r}(\mathfrak{n}, \mathfrak{h}))$ . The intuition in Example 5.1 will be formulated properly in Section 9.1, but we shall first deal with more general multisegments (rather than only two segments) below.

5.2. Local minimizability. We now define minimizability in Definition 5.2 to show the uniqueness for the  $\leq_Z$ -minimal element in  $\mathcal{S}(\pi, \tau)$  in Theorem 6.4.

**Definition 5.2.** Let  $\mathfrak{h} \in \text{Mult}_{\rho}$  and let  $\mathfrak{n} \in \text{Mult}_{\rho}$  be admissible to  $\mathfrak{h}$ . Let a be the smallest integer such that  $\mathfrak{n}[a] \neq \emptyset$ . We say that  $(\mathfrak{n}, \mathfrak{h})$  is *locally minimizable* if there exists a segment  $\overline{\Delta}$  in  $\mathfrak{n}[a+1]$  such that the following holds:

 $|\left\{\Delta \in \mathfrak{n}[a] : \overline{\Delta} \subset \Delta\right\}| < |\left\{\Delta \in \mathfrak{fs}(\mathfrak{n}\mathfrak{h}) : \overline{\Delta} \subset \Delta\right\}|.$ 

We emphasis that the non-strict inequality  $\leq$  always holds.

**Remark 5.3.** We give more explanations on Definition 5.2. As suggested from the terminology, those locally minimizable  $(\mathfrak{n}, \mathfrak{h})$  is to find some  $\mathfrak{n}' <_Z \mathfrak{n}$  such that  $\mathfrak{r}(\mathfrak{n}', \mathfrak{h}) = \mathfrak{r}(\mathfrak{n}, \mathfrak{h})$ . For instance, if all segments  $\Delta$  in  $\mathfrak{n}[a]$  satisfy  $\overline{\Delta} \subset \Delta$ , the removal process guarantees that any  $\Delta$  in  $\mathfrak{fs}(\mathfrak{n}, \mathfrak{h})$  also satisfies  $\overline{\Delta} \subset \Delta$ . Hence, the inequality in Definition 5.2 is not satisfied. On the other hand, all segments in  $\mathfrak{n}[a]$  are not linked to  $\overline{\Delta}$  and so there is no intersection-union operation for segments in  $\mathfrak{n}[a]$  and  $\overline{\Delta}$ .

We have one simple way to check local minimizable by just working on one segment. The proof is straightforward from definitions:

**Lemma 5.4.** Let  $\mathfrak{h} \in \operatorname{Mult}_{\rho}$  and let  $\mathfrak{n} \in \operatorname{Mult}_{\rho}$  be admissible to  $\mathfrak{h}$ . Let a be the smallest integer such that  $\mathfrak{n}[a] \neq \emptyset$ . Let  $\Delta$  be a segment in  $\mathfrak{n}[a]$ . If there exists a segment  $\overline{\Delta}$  in  $\mathfrak{n}[a+1]$  such that  $\overline{\Delta} \not\subset \Delta$  and  $\overline{\Delta} \subset \Upsilon(\Delta, \mathfrak{h})$ , then  $(\mathfrak{n}, \mathfrak{h})$  is not locally minimizable.

**Example 5.5.** Let  $\mathfrak{h} = \{[0,1]_{\rho}, [1,4]_{\rho}, [1,5]_{\rho}, [1,6]_{\rho}, [2,5]_{\rho}, [3,4]_{\rho}\}, \text{let } \mathfrak{n} = \{[1,3]_{\rho}, [1,6]_{\rho}, [2,4]_{\rho}\}$  and let  $\mathfrak{n}' = \{[1,3]_{\rho}, [1,6]_{\rho}, [2,5]_{\rho}\}.$ 



The blue points represent  $\mathfrak{fs}(\mathfrak{n},\mathfrak{h})$  and  $\mathfrak{fs}(\mathfrak{n}',\mathfrak{h})$ . Note that

0

$$|\{\Delta \in \mathfrak{n}[1] : [2,4]_{\rho} \subset \Delta\}| = 1, \quad |\{\Delta \in \mathfrak{fs}(\mathfrak{n},\mathfrak{h}) : [2,4]_{\rho} \subset \Delta\}| = 2$$

and so (n, h) is locally minimizable. On the other hand,

 $|\{\Delta \in \mathfrak{n}'[1] : [2,5]_{\rho} \subset \Delta\}| = 1, \quad |\{\Delta \in \mathfrak{fs}(\mathfrak{n}',\mathfrak{h}) : [2,5]_{\rho} \subset \Delta\}| = 1.$ 

Hence  $(\mathfrak{n}', \mathfrak{h})$  is not locally minimizable.

**Lemma 5.6.** Let  $\mathfrak{h} \in \operatorname{Mult}_{\rho}$  and let  $\mathfrak{n} \in \operatorname{Mult}_{\rho}$  be admissible to  $\mathfrak{h}$ . Let  $\mathfrak{n}' = \mathfrak{trd}(\mathfrak{n}, \mathfrak{h})$  and  $\mathfrak{h}' = \mathfrak{trr}(\mathfrak{n},\mathfrak{h})$ . Let a be the smallest integer such that  $\mathfrak{n}'[a] \neq \emptyset$ . Fix some c > a + 1. Fix a segment  $\overline{\Delta}$  of the form  $[c,d]_{\rho}$  for some d. Suppose

$$(*) \quad |\left\{\Delta \in \mathfrak{n}' : \overline{\Delta} \subset \Delta\right\}| < |\left\{\Delta \in \mathfrak{fs}(\mathfrak{n}', \mathfrak{h}') : \overline{\Delta} \subset \Delta\right\}|.$$

• There exists a segment  $\widetilde{\Delta}$  in  $\mathfrak{n}[a] + \mathfrak{n}[a+1]$  such that

$$\mathfrak{fs}(\mathfrak{n},\mathfrak{h})=\mathfrak{fs}(\widetilde{\mathfrak{n}},\mathfrak{h}),\quad\mathfrak{fs}(\mathfrak{n}',\mathfrak{h}')=\mathfrak{fs}(\widetilde{\mathfrak{n}}',\mathfrak{h}'),$$

where  $\widetilde{\mathfrak{n}}$  is obtained from  $\mathfrak{n}$  by an elementary intersection-union process between  $\widetilde{\Delta}$ and  $\overline{\Delta}$ , and  $\widetilde{\mathfrak{n}}' = \mathfrak{trd}(\mathfrak{n}, \mathfrak{h})$ .

• Furthermore, if the segment  $\widetilde{\Delta}$  cannot be chosen in  $\mathfrak{n}[a+1]$ , then  $|\{\Delta \in \mathfrak{n}[a] : \overline{\Delta} \subset \Delta\}| < \mathbb{N}$  $|\left\{\Delta \in \mathfrak{fs}(\mathfrak{n},\mathfrak{h}): \overline{\Delta} \subset \Delta\right\}|.$ 

*Proof.* Recall that

$$\mathfrak{l}(\mathfrak{n}[a]) = \left\{ {}^{-}\Delta : \Delta \in \mathfrak{n}[a], \Delta \neq [a] 
ight\}.$$

Let  $r = |{}^{-}(\mathfrak{n}[a])|$  and let  $s = |\mathfrak{n}[a+1]|$ . We arrange the segments in  $\mathfrak{n}'$  as follows: the first r segments are those in  $(\mathfrak{n}[a])$ , and the remaining segments are those in  $\mathfrak{n}[a+1]$ . To facilitate discussions, the first r segments are labelled as

$$\Delta_1,\ldots,\Delta_r$$

and the remaining segments are:

$$\widetilde{\Delta}_1, \ldots, \widetilde{\Delta}_s$$

**.** .

We also set

$$\Lambda_{i} = \Upsilon(\Delta_{i}, \mathfrak{r}(\{\Delta_{1}, \dots, \Delta_{i-1}\}, \mathfrak{h}'),$$
$$\widetilde{\Lambda}_{i} = \Upsilon(\widetilde{\Delta}_{i}, \mathfrak{r}(\{\widetilde{\Delta}_{1}, \dots, \widetilde{\Delta}_{i-1}, \Delta_{1}, \dots, \Delta_{r}\}, \mathfrak{h}').$$

Case 1:

$$|\left\{\Delta \in \mathfrak{n}[a] : \overline{\Delta} \subset \Delta\right\}| = |\left\{\Delta \in \mathfrak{fs}(\mathfrak{n},\mathfrak{h}) : \overline{\Delta} \subset \Delta\right\}|.$$

This condition and the nesting property implies that for the first r segments  $\Delta_i$ , if  $\overline{\Delta} \not\subset \Delta_i$ , then

 $\overline{\Delta} \not\subset \Lambda_i.$ 

Thus condition (\*) implies that there exists a segment  $\widetilde{\Delta}_i$  such that  $\overline{\Delta} \not\subset \widetilde{\Delta}_i$  and  $\overline{\Delta} \subset \Lambda_i$ . Now we do the intersection-union operation on  $\overline{\Delta}$  and  $\widetilde{\Delta}_i$  to obtain  $\widetilde{\mathfrak{n}}$  from  $\mathfrak{n}$ . Then  $\mathfrak{n}[a] = \widetilde{\mathfrak{n}}[a]$  and so  $\mathfrak{fs}(\mathfrak{n},\mathfrak{h}) = \mathfrak{fs}(\widetilde{\mathfrak{n}},\mathfrak{h})$ . And,  $\mathfrak{n}'$  and  $\widetilde{\mathfrak{n}}' := \mathfrak{trd}(\widetilde{\mathfrak{n}},\mathfrak{h}')$  are only differed by  $\widetilde{\Delta}_i$  and  $\widetilde{\Delta}_i \cup \overline{\Delta}$ . However, if we impose the same ordering in computing  $\mathfrak{fs}(\widetilde{\mathfrak{n}}',\mathfrak{h}')$ , it is straightforward to use  $\overline{\Delta} \subset \Lambda_i$  to see that

$$\mathfrak{fs}(\mathfrak{n}',\mathfrak{h}')=\mathfrak{fs}(\widetilde{\mathfrak{n}}',\mathfrak{h}').$$

Case 2:

$$\left\{\Delta \in \mathfrak{n}[a] : \overline{\Delta} \subset \Delta\right\} | < |\left\{\Delta \in \mathfrak{fs}(\mathfrak{n},\mathfrak{h}) : \overline{\Delta} \subset \Delta\right\} |.$$

Now, by (\*), there exists a segment  $\widetilde{\Delta} = \Delta_i$  or  $\widetilde{\Delta}_i$  in  $\mathfrak{n}'$  such that  $\overline{\Delta} \not\subset \widetilde{\Delta}$  and  $\overline{\Delta} \subset \Lambda$ , where  $\Lambda = \Lambda_i$  or  $\widetilde{\Lambda}_i$  according to  $\widetilde{\Delta}$ .

If  $\overline{\Delta} = \overline{\Delta}_i$  for some *i*, then the intersection-union operation is done between  $\overline{\Delta}$  and  $\overline{\Delta}$ . The argument is similar to Case 1 and we omit the details.

We now consider the case that  $\Delta = \Delta_i$  for some *i*. For convenience, set  $+[a+1,c]_{\rho} = [a,c]_{\rho}$  for any *c*. Note that all  $+\Delta_k$   $(k = 1, \ldots, r)$  constitute all the non-singleton segments sin  $\mathfrak{n}[a]$ . We can use the ordering  $+\Delta_1, \ldots, +\Delta_r$  (with other singleton segments at the end) to compute  $\mathfrak{fs}(\mathfrak{n},\mathfrak{h})$ ; and similarly use that ordering with  $+\Delta_i$  replaced by  $+\Delta_i \cup \overline{\Delta}$  to compute  $\mathfrak{fs}(\mathfrak{n},\mathfrak{h})$ . The only difference is to compute the first segments for  $+\Delta_i$  and  $+\Delta_i \cup \overline{\Delta}$ , but we can still guarantee that choices for first segments (for computing  $\mathfrak{fs}(\mathfrak{n},\mathfrak{h})$  and  $\mathfrak{fs}(\mathfrak{n}',\mathfrak{h}')$ ) still coincide by using the nesting property of the removal process and the condition  $\overline{\Delta} \subset \Lambda$ . Hence,  $\mathfrak{fs}(\mathfrak{n},\mathfrak{h}) = \mathfrak{fs}(\mathfrak{\tilde{n}},\mathfrak{h})$ . Computing  $\mathfrak{fs}(\mathfrak{n}',\mathfrak{h}') = \mathfrak{fs}(\mathfrak{\tilde{n}}',\mathfrak{h}')$  is again similar since the only difference between  $\mathfrak{n}'$  and  $\mathfrak{\tilde{n}}'$  is  $\Delta_i$  and  $\Delta_i \cup \overline{\Delta}$ .

#### 6. Uniqueness of minimality in $S(\pi, \tau)$

The terminology of minimizability is suggested by the following lemma:

**Lemma 6.1.** Let  $\mathfrak{h} \in \text{Mult}_{\rho}$  and let  $\mathfrak{n} \in \text{Mult}_{\rho}$  be admissible to  $\mathfrak{h}$ . Let the fine chain  $\text{fc}_{\mathfrak{h}}(\mathfrak{n})$  take the form

$$\mathfrak{fs}(\mathfrak{n}_0,\mathfrak{h}_0),\mathfrak{fs}(\mathfrak{n}_1,\mathfrak{h}_1),\ldots$$

as in Definition 3.6. If  $(\mathfrak{n}_j,\mathfrak{h}_j)$  is not locally minimizable for any j, then there is no multisegment  $\mathfrak{n}'$  such that  $\mathfrak{n}' \leq_Z \mathfrak{n}$  and  $\mathfrak{r}(\mathfrak{n},\mathfrak{h}) = \mathfrak{r}(\mathfrak{n}',\mathfrak{h})$ .

*Proof.* Suppose there is a multisegment  $\mathfrak{n}'$  such that  $\mathfrak{n}' \leq_Z \mathfrak{n}$  and  $\mathfrak{r}(\mathfrak{n}, \mathfrak{h}) = \mathfrak{r}(\mathfrak{n}', \mathfrak{h})$ . Then, by Theorem 4.4, we may take  $\mathfrak{n}'$  to be obtained from  $\mathfrak{n}$  by an elementary intersection-union process. Let  $\widetilde{\Delta} = [\widetilde{a}, \widetilde{b}]_{\rho}, \overline{\Delta} = [\overline{a}, \overline{b}]_{\rho}$  in  $\mathfrak{n}$  be the segments involved in the intersection-union process, and switching labeling if necessary, we may assume that  $\widetilde{a} < \overline{a}$ .

We similarly obtain the fine chain

$$\mathfrak{fs}(\mathfrak{n}_0',\mathfrak{h}_0'),\mathfrak{fs}(\mathfrak{n}_1',\mathfrak{h}_1'),\ldots,$$

for  $(\mathfrak{n}', \mathfrak{h})$ . We consider j such that  $\overline{a} - 1$  is the smallest integer c such that  $\mathfrak{n}_j[c] \neq \emptyset$ . (Such j exists by using the condition that  $\widetilde{\Delta}$  and  $\overline{\Delta}$  are linked.)

Then, in  $\mathfrak{n}_j$ , we have a segment  $[\overline{a}-1, \widetilde{b}]_{\rho}$  coming by truncating  $\widetilde{\Delta}$ . If we replace  $[\overline{a}-1, \widetilde{b}]_{\rho}$  in  $\mathfrak{n}_j$  by  $[\overline{a}-1, \overline{b}]_{\rho}$ , this gives  $\mathfrak{n}'_j$ .

Now

$$\left|\left\{\Delta \in \mathfrak{n}_{j}: \widetilde{\Delta} \subset \Delta\right\}\right| < \left|\left\{\Delta \in \mathfrak{n}_{j}': \widetilde{\Delta} \subset \Delta\right\}\right| \le \left|\left\{\Delta \in \mathfrak{fs}(\mathfrak{n}_{j}', \mathfrak{h}_{j}): \widetilde{\Delta} \subset \Delta\right\}\right|,$$

where the first strict inequality comes from  $[\overline{a} - 1, \overline{b}]_{\rho}$ . But by Lemma 3.9, two fine chains coincide and in particular  $\mathfrak{fs}(\mathfrak{n}_j, \mathfrak{h}_j) = \mathfrak{fs}(\mathfrak{n}'_j, \mathfrak{h}_j)$ . Hence, we now have that  $(\mathfrak{n}_j, \mathfrak{h}_j)$  is locally minimizable by Definition 5.2 as desired.

We now prove the converse of Lemma 6.1. The main idea is to use Lemma 5.6 to locate a suitable choice of a segment for the intersection-union process. Since the local minimizability is for the multisegments in the fine chain  $fc_{\mathfrak{h}}(\mathfrak{n})$ , one may not be able to immediately find a segment that originally comes from  $\mathfrak{n}$  and so the second bullet of Lemma 5.6 allows one to trace back and inductively use Lemma 5.6 to find suitable segments coming from an original segment in  $\mathfrak{n}$  responsible for the intersection-union process.

**Lemma 6.2.** We keep using the notations in Lemma 6.1. If  $(\mathfrak{n}_j, \mathfrak{h}_j)$  is locally minimizable for some j, then there is a multisegment  $\mathfrak{n}'$  such that  $\mathfrak{n}' \leq_Z \mathfrak{n}$  and  $\mathfrak{r}(\mathfrak{n}, \mathfrak{h}) = \mathfrak{r}(\mathfrak{n}', \mathfrak{h}')$ .

*Proof.* We pick any j such that  $(\mathfrak{n}_j, \mathfrak{h}_j)$  is locally minimizable. Let a be the smallest integer such that  $\mathfrak{n}_j[a] \neq \emptyset$ . The below argument is similar if j = 1 and so we assume j > 1 for convenience of the stated form of Lemma 5.6.

Note that  $\mathfrak{n}_j[a] = -(\mathfrak{n}_{j-1}[a-1]) + \mathfrak{n}_j[a]$ . (Here we have  $\mathfrak{n}_{j-1}[a] = \mathfrak{n}[a]$ .) The local minimizability condition implies that we can use the first bullet of Lemma 5.6 (set c = a+1 in our case) with respect to a certain segment in  $\mathfrak{n}[a+1]$ , denoted by  $\overline{\Delta}$ . Then, that lemma implies that we can find a segment  $\widetilde{\Delta}$  in  $-(\mathfrak{n}_{j-1}[a-1]) + \mathfrak{n}[a]$  satisfying the required properties in the lemma.

The first case is that  $\overline{\Delta}$  comes from  $\mathfrak{n}[a]$ . In this case, let  $\widetilde{\mathfrak{n}}$  be the multisegment obtained from  $\mathfrak{n}$  by the intersection-union operation of the segments  $\overline{\Delta}$  and  $\overline{\Delta}$ . Then it is straightforward from definitions that

$$\mathfrak{fs}(\mathfrak{n}_0,\mathfrak{h}_0)=\mathfrak{fs}(\widetilde{\mathfrak{n}}_0,\mathfrak{h}_0),\ldots,\mathfrak{fs}(\mathfrak{n}_{j-1},\mathfrak{h}_{j-1})=\mathfrak{fs}(\widetilde{\mathfrak{n}}_{j-1},\mathfrak{h}_{j-1}),$$

where  $\mathfrak{h}_0 = \mathfrak{h}$ ,  $\mathfrak{fs}(\tilde{\mathfrak{n}}_k, \mathfrak{h}_k)$  are the first j-1 terms of  $\mathrm{fc}(\tilde{\mathfrak{n}}, \mathfrak{h})$ . However,  $\mathfrak{fs}(\mathfrak{n}_j, \mathfrak{h}_j) = \mathfrak{fs}(\tilde{\mathfrak{n}}_j, \mathfrak{h}_j)$  is guaranted by Lemma 5.6. But then  $\mathfrak{n}_{j+1} = \tilde{\mathfrak{n}}_{j+1}$  and so the reamining terms in two fine chains also agree. Hence, two fine chains coincide and so  $\mathfrak{r}(\mathfrak{n}, \mathfrak{h}) = \mathfrak{r}(\tilde{\mathfrak{n}}, \mathfrak{h})$  by Lemma 3.9.

The second case is that  $\Delta$  cannot come from  $\mathfrak{n}[a]$ . In such case, the second bullet of Lemma 5.6 implies that we have

$$|\left\{\Delta \in \mathfrak{n}_{j-1}[a-1]: \overline{\Delta} \subset \Delta\right\}| < |\left\{\Delta \in \mathfrak{fs}(\mathfrak{n}_{j-1}[a-1], \mathfrak{h}): \overline{\Delta} \subset \Delta\right\}|$$

But then, we can apply Lemma 5.6 again to find another segment  $\widetilde{\Delta}$ . If such segment  $\widetilde{\Delta}$  can be found in  $(\mathfrak{n}_{j-2}[a-1])$ , then we repeat the similar argument of the first case above. Otherwise, we apply Lemma 5.6 again. In those cases, the coincidence of the fine chains are guaranteed by Lemma 5.6. Hence, we also have  $\mathfrak{r}(\mathfrak{n},\mathfrak{h}) = \mathfrak{r}(\widetilde{\mathfrak{n}},\mathfrak{h})$  by Lemma 3.9. (Strictly speaking in terms of the way in stating Lemma 5.6, one has to trace the proof to see that the choices for  $\widetilde{\Delta}$  and  $\widetilde{\widetilde{\Delta}}$  in each step can be made to agree after truncating the point  $[a-1]_{\rho}$ . Since there is no new idea on that, we avoid further notation complications.)  $\Box$ 

We explain the main idea of the proof for Proposition 6.3 below, which is inductive in nature. One first picks two minimal multisegments  $\mathfrak{n}$  and  $\mathfrak{n}'$  in  $\mathcal{S}'(\mathfrak{h},\mathfrak{p})$ . One then finds  $\prec^{L}$ -minimal segments  $\widetilde{\Delta}$  and  $\widetilde{\Delta}'$  in  $\mathfrak{n}$  and  $\mathfrak{n}'$  respectively. If  $\widetilde{\Delta} = \widetilde{\Delta}'$ , then one uses induction to argue  $\mathfrak{n} - \widetilde{\Delta} = \mathfrak{n}' - \widetilde{\Delta}$ . If  $\widetilde{\Delta} \neq \widetilde{\Delta}'$ , then one first reduces to the case that  $\widetilde{\Delta} \subsetneq \widetilde{\Delta}'$ . Then one applies induction hypothesis to show  $\widetilde{\Delta}'$  is also in  $\mathfrak{n}$ . Then one shows that  $\widetilde{\Delta}$  and  $\widetilde{\Delta}'$ in  $\mathfrak{n}$  give rise the local minimizability.

**Proposition 6.3.** Let  $\mathfrak{h} \in \text{Mult}_{\rho}$ . Then there exists a unique minimal element in  $\mathcal{S}'(\mathfrak{h}, \mathfrak{p})$  if  $\mathcal{S}'(\mathfrak{h}, \mathfrak{p}) \neq \emptyset$ .

*Proof.* Let  $\mathfrak{n}, \mathfrak{n}'$  be two minimal multisegments in  $\mathcal{S}'(\mathfrak{h}, \mathfrak{p})$ . Let a be the smallest integer such that  $\mathfrak{n}[a] \neq 0$ . Then, by a comparison on cuspidal representations, a is also the smallest integer such that  $\mathfrak{n}'[a] \neq 0$ .

Suppose  $\mathfrak{n}[a] \cap \mathfrak{n}'[a] \neq \emptyset$ . Let  $\widetilde{\Delta} \in \mathfrak{n}[a] \cap \mathfrak{n}'[a]$ . Then we consider  $\mathcal{S}'(\mathfrak{r}(\widetilde{\Delta},\mathfrak{h}),\mathfrak{p})$ . The minimality for  $\mathfrak{n}$  and  $\mathfrak{n}'$  also guarantees that  $\mathfrak{n} - \widetilde{\Delta}$  and  $\mathfrak{n}' - \widetilde{\Delta}$  are also minimal in  $\mathcal{S}'(\mathfrak{r}(\widetilde{\Delta},\mathfrak{h}),\mathfrak{p})$ . Thus, by induction, we have that  $\mathfrak{n} - \widetilde{\Delta} = \mathfrak{n}' - \widetilde{\Delta}$  and so  $\mathfrak{n} = \mathfrak{n}'$ .

Now suppose  $\mathfrak{n}[a] \cap \mathfrak{n}'[a] = \emptyset$  to obtain a contradiction. Let  $\Delta$  and  $\Delta'$  be the shortest segment in  $\mathfrak{n}[a]$  and  $\mathfrak{n}'[a]$  respectively. Switching labeling if necessary, we may assume that  $\Delta \subsetneq \Delta'$ . Then, by Lemma 3.4,

$$\mathfrak{r}(\mathfrak{n},\mathfrak{h}) = \mathfrak{r}(\mathfrak{trd}(\mathfrak{n},\mathfrak{h}),\mathfrak{trr}(\mathfrak{n},\mathfrak{h})), \quad \mathfrak{r}(\mathfrak{n}',\mathfrak{h}) = \mathfrak{r}(\mathfrak{trd}(\mathfrak{n}',\mathfrak{h}),\mathfrak{trr}(\mathfrak{n}',\mathfrak{h})).$$

By Lemma 3.9,

$$\operatorname{trr}(\mathfrak{n},\mathfrak{h}) = \operatorname{trr}(\mathfrak{n}',\mathfrak{h}).$$

By Lemma 6.2,  $(\mathfrak{trr}(\mathfrak{n},\mathfrak{h}),\mathfrak{trd}(\mathfrak{n},\mathfrak{h}))$  and the terms from the fine chains are not locally minimizable. Similarly, this also holds for  $(\mathfrak{trr}(\mathfrak{n}',\mathfrak{h}),\mathfrak{trd}(\mathfrak{n},\mathfrak{h}))$ . However, Lemma 6.1 implies that both  $\mathfrak{trd}(\mathfrak{n},\mathfrak{h})$  and  $\mathfrak{trd}(\mathfrak{n}',\mathfrak{h})$  are minimal in  $\mathcal{S}'(\mathfrak{trr}(\mathfrak{n},\mathfrak{h}),\mathfrak{p}) = \mathcal{S}'(\mathfrak{trr}(\mathfrak{n}',\mathfrak{h}),\mathfrak{p})$ . Hence, by induction,

$$\mathfrak{trd}(\mathfrak{n},\mathfrak{h}) = \mathfrak{trd}(\mathfrak{n}',\mathfrak{h}).$$

But then, the disjointness assumption implies that  $^{-}\Delta' \in \mathfrak{n}$ . But  $^{-}\Delta' \not\subset \Delta$  and  $^{-}\Delta' \subset \Upsilon(\Delta, \mathfrak{h})$ . By Lemma 5.4, this implies

$$\left|\left\{\widetilde{\Delta}\in\mathfrak{n}[a]: {}^{-}\Delta'\subset\widetilde{\Delta}\right\}\right|<\left|\left\{\widetilde{\Delta}\in\mathfrak{fs}(\mathfrak{n},\mathfrak{h}): {}^{-}\Delta'\subset\widetilde{\Delta}\right\}\right|.$$

Hence, (n, b) is locally minimizable. This contradicts to Lemma 6.2.

**Theorem 6.4.** Let  $\pi \in \operatorname{Irr}_{\rho}$  and let  $\tau$  be a simple quotient of  $\pi^{(i)}$  for some *i*. Then  $\mathcal{S}(\pi, \tau)$  has a unique minimal element if  $\mathcal{S}(\pi, \tau) \neq \emptyset$ . Here the minimality is with respect to  $\leq_Z$ .

*Proof.* This follows from Proposition 6.3 and Theorem 2.4.

#### 7. Examples of minimality

#### 7.1. Minimality for the highest derivative multisegment.

**Theorem 7.1.** Let  $\pi \in \operatorname{Irr}_{\rho}$ . Then  $\mathfrak{ho}(\pi)$  is minimal in  $\mathcal{S}(\pi, \pi^{-})$ .

*Proof.* It is shown in [Ch25] that  $D_{\mathfrak{hd}(\pi)}(\pi) \cong \pi^-$ . It remains to prove that  $\mathfrak{hd}(\pi)$  is minimal in  $\mathcal{S}(\pi,\pi^-)$ . Theorem 4.4 reduces to show that if  $\mathfrak{n}$  is a multisegment obtained by an elementary intersection-union process from  $\mathfrak{hd}(\pi)$ , then  $D_{\mathfrak{n}}(\pi) = 0$ .

Let  $\Delta_1 = [a_1, b_1]_{\rho}, \Delta_2 = [a_2, b_2]_{\rho}$  be two linked segments in  $\mathfrak{ho}(\pi)$ . Relabeling if necessary, we assume that  $a_1 < a_2$ . Define

$$\mathfrak{n} = \mathfrak{h}\mathfrak{d}(\pi) - \{\Delta_1, \Delta_2\} + \Delta_1 \cup \Delta_2 + \Delta_1 \cap \Delta_2.$$

14

Then,  $\mathfrak{n}[e] = \mathfrak{ho}(\pi)[e]$  for any  $e < a_1$  and  $\mathfrak{n}[a_1] \not\leq_{a_1} \mathfrak{ho}(\pi)[a_1]$ . Hence, by Theorems 2.4 and 2.5,

$$D_{\mathfrak{n}[a_1]} \dots D_{\mathfrak{n}[c]}(\pi) = D_{\mathfrak{n}[a_1]} \circ D_{\mathfrak{h}\mathfrak{d}(\pi)[a_1-1]} \circ \dots \circ D_{\mathfrak{h}\mathfrak{d}(\pi)[c]}(\pi) = 0,$$

and so  $\mathfrak{n} \notin \mathcal{S}(\pi, \pi^{-})$ . Here c is the smallest integer such that  $\mathfrak{ho}(\pi)[c] \neq 0$ .

### 7.2. Minimal multisegment for the generic case.

**Proposition 7.2.** Let  $\pi \in \operatorname{Irr}_{\rho}$  be generic. Let  $\tau$  be a (generic) simple quotient of  $\pi^{(i)}$  for some *i*. Then the minimal multisegment in  $S(\pi, \tau)$  is generic *i.e.* any two segments in the minimal multisegment are unlinked.

One may prove the above proposition by some analysis of derivative resultant multisegments. We shall give another proof using the following lemma:

**Lemma 7.3.** Let  $\pi \in \operatorname{Irr}_{\rho}(G_n)$  be generic. For any *i*, and for any irreducible submodule  $\tau_1 \boxtimes \tau_2$  of  $\pi_{N_i}$  as  $G_{n-i} \times G_i$ -representations, both  $\tau_1$  and  $\tau_2$  are generic.

*Proof.* Recall that for a generic  $\pi$ ,

$$\pi \cong \operatorname{St}(\Delta_1) \times \ldots \times \operatorname{St}(\Delta_r)$$

for mutually unlinked segments  $\Delta_1, \ldots, \Delta_r$ .

Now, with a suitable arrangement on the orderings of the segments, one may argue as in [Ch21, Corollary 2.6] to have that a simple quotient of  $\pi_{N_i}$  takes the form  $\tau \boxtimes \omega$  for some generic  $\tau \in \operatorname{Irr}(G_{n-i})$ . Hence it remains to show  $\omega$  is also generic. We consider

$$\pi_{N_i} \twoheadrightarrow \tau \boxtimes \omega$$

and taking the twisted Jacquet functor on the  $G_{n-i}$ -parts yields that

$$^{(n-i)}\pi \twoheadrightarrow \omega.$$

Now using [Ch21, Corollary 2.6] for left derivatives, we have that  $\omega$  is also generic as desired.

Proof of Proposition 7.2. Let  $\pi \in \operatorname{Irr}_{\rho}$  be generic and let  $\tau$  be a simple (generic) quotient of  $\pi^{(i)}$  (see [Ch21, Corollary 2.6]). Then,  $\pi_{N_i}$  has a simple quotient of the form  $\tau \boxtimes \omega$  for some  $\omega \in \operatorname{Irr}_{\rho}(G_i)$ . By Lemma 7.3,  $\omega$  is also generic and hence  $\omega \cong \operatorname{St}(\Delta_1) \times \ldots \times \operatorname{St}(\Delta_k)$ for some mutually unlinked segments  $\Delta_1, \ldots, \Delta_k$ . Now,  $\pi$  is the unique submodule of  $\tau \times \operatorname{St}(\Delta_1) \times \ldots \times \operatorname{St}(\Delta_k)$ . By a standard argument, we have that:

$$D_{\Delta_1} \circ \ldots \circ D_{\Delta_k}(\pi) \cong \tau.$$

Hence,  $\{\Delta_1, \ldots, \Delta_k\} \in \mathcal{S}(\pi, \tau)$ . The minimality of  $\{\Delta_1, \ldots, \Delta_k\}$  is automatic since any generic multisegment is minimal in  $\operatorname{Mult}_{\rho}$  with respect to  $\leq_Z$ . Now the statement follows from the uniqueness in Theorem 6.4.

### 8. Non-uniqueness of maximal elements in $S(\pi, \tau)$

8.1. Highest derivative multisegments. Let  $\pi \in \operatorname{Irr}_{\rho}$ . Then  $\mathcal{S}(\pi, \pi^{-})$  contains a unique maximal multisegment, and such multisegment has all segments to be singletons. Combining with Theorem 4.4, one can describe all multisegments in  $\mathcal{S}(\pi, \pi^{-})$ .

8.2. Failure of uniqueness of maximality. As mentioned in [Ch25], in general, derivatives of cuspidal representations are not enough for constructing all simple quotients of Bernstein-Zelevinsky derivatives, and the set  $S(\pi, \tau)$  may contain some multisegments whose segments are not all singletons. We give an example to show that in general, there is no uniqueness for  $\leq_Z$ -maximal elements in  $S(\pi, \tau)$ .

 $\operatorname{Let}$ 

$$\mathfrak{h} = \{ [0,3]_{\rho}, [0,1]_{\rho}, [1,2]_{\rho}, [1,2]_{\rho}, [2]_{\rho}, [3]_{\rho} \}.$$

Let  $\mathfrak{n} = \{[0,3]_{\rho}, [1,2]_{\rho}\}$ . Then

$$\mathfrak{r} := \mathfrak{r}(\mathfrak{n}, \mathfrak{h}) = \{ [0, 1]_{\rho}, [1, 2]_{\rho}, [2]_{\rho}, [3]_{\rho} \} = \mathfrak{h} - \mathfrak{n}.$$

We claim that

$$\mathcal{S}'(\mathfrak{h},\mathfrak{r}) = \{\mathfrak{n}, \{[0,3]_{\rho}, [1]_{\rho}, [2]_{\rho}\}, \{[0,2]_{\rho}, [1,3]_{\rho}\}\}.$$

It is direct to check that the three elements are in  $\mathcal{S}(\mathfrak{h},\mathfrak{r})$ , and the last two elements are both maximal.

To see that there are no more elements, we first observe that any multisegment  $\mathfrak{n}'$  in  $\mathcal{S}'(\mathfrak{h},\mathfrak{r})$  has only one segment  $\widetilde{\Delta}$  with  $a(\widetilde{\Delta}) = \nu^0$ . By considering the first segment in the removal sequence  $\mathfrak{r}(\widetilde{\Delta},\mathfrak{h})$ , we note that  $[0]_{\rho}, [0,1]_{\rho} \notin \mathcal{S}'(\mathfrak{h},\mathfrak{r})$ . In other words,  $[0,2]_{\rho}$  or  $[0,3]_{\rho}$  in  $\mathcal{S}'(\mathfrak{h},\mathfrak{r})$ . It remains to check that the following three elements:

$$\{[0,2]_{\rho}, [1]_{\rho}, [2]_{\rho}, [3]_{\rho}\}, \{[0,2]_{\rho}, [1,2]_{\rho}, [3]_{\rho}\}, \{[0,2]_{\rho}, [1]_{\rho}, [2,3]_{\rho}\}$$

are not in  $\mathcal{S}'(\mathfrak{h}, \mathfrak{r})$ , which is straightforward.

To show that the maximality for  $S(\pi, \tau)$  is not unique in general, one still needs to ask whether there exists  $\pi \in \operatorname{Irr}_{\rho}$  such that  $\mathfrak{ho}(\pi) = \mathfrak{h}$ . This is indeed the case and we shall postpone proving this in the sequel [Ch24+e], where we shall need such fact in a more substantial way.

### 9. MINIMALITY FOR TWO SEGMENT CASE

In this section, we study the minimality for two segment cases.

### 9.1. Non-overlapping property.

**Definition 9.1.** Let  $\mathfrak{h} \in \operatorname{Mult}_{\rho}$ . Let  $\Delta \in \operatorname{Seg}_{\rho}$  be admissible to  $\mathfrak{h}$ . Let  $\Delta' \in \operatorname{Seg}_{\rho}$  linked to  $\Delta$  with  $\Delta' > \Delta$ . We say that the triple  $(\Delta, \Delta', \mathfrak{h})$  satisfies the *non-overlapping property* if for the shortest segment  $\overline{\Delta}$  in the removal sequence for  $(\Delta, \mathfrak{h})$  that contains  $\nu^{-1}a(\Delta')$ , we have  $\Delta' \not\subset \overline{\Delta}$ . (We remark that for later applications, we do not impose the condition that  $\Delta'$  is admissible to  $\mathfrak{h}$ .)

**Example 9.2.** (1) Let  $\mathfrak{h} = \{[0,7]_{\rho}, [3,6]_{\rho}, [6,10]_{\rho}\}$ . Let  $\Delta = [0,5]_{\rho}$  and let  $\Delta' = [6,7]_{\rho}$ . Then  $(\Delta, \Delta', \mathfrak{h})$  satisfies the non-overlapping property.



The blue points are those points removed by applying  $\mathfrak{r}(\Delta, .)$  while the red points are those points removed by applying  $\mathfrak{r}(\Delta', .)$ . Note that the shortest segment in the removal sequence containing  $[5]_{\rho}$  is  $[3, 6]_{\rho}$ , which does not contain  $[6, 7]_{\rho}$ .

(2) Let  $\mathfrak{h} = \{[0, 8]_{\rho}, [3, 6]_{\rho}, [6, 10]_{\rho}\}$ . Let  $\Delta = [0, 7]_{\rho}$  and let  $\Delta' = [6, 8]_{\rho}$ . Then  $(\Delta, \Delta', \mathfrak{h})$  does not satisfy the non-overlapping property. The graph for carrying out the removal sequence looks like:



In the graph above, the segment  $[0, 8]_{\rho}$  contains  $[5]_{\rho}$ , and  $[6, 8]_{\rho} \subset [0, 8]_{\rho}$ .

**Lemma 9.3.** Let  $\mathfrak{h} \in \text{Mult}_{\rho}$ . Let  $\Delta \in \text{Seg}_{\rho}$  be admissible to  $\mathfrak{h}$ . Let  $\Delta' \in \text{Seg}_{\rho}$  be admissible to  $\mathfrak{r}(\Delta, \mathfrak{h})$ . Suppose  $\Delta'$  is linked to  $\Delta$  with  $\Delta' > \Delta$ . Then  $(\Delta, \Delta', \mathfrak{h})$  does not satisfy the non-overlapping property if and only if

$$\mathfrak{r}(\{\Delta \cap \Delta', \Delta \cup \Delta'\}, \mathfrak{h}) = \mathfrak{r}(\{\Delta, \Delta'\}, \mathfrak{h}).$$

*Proof.* Let  $\mathfrak{n} = \{\Delta, \Delta'\}$ . Suppose  $\mathfrak{r}(\{\Delta \cap \Delta', \Delta \cup \Delta'\}, \mathfrak{h}) \neq \mathfrak{r}(\{\Delta, \Delta'\}, \mathfrak{h})$ . Lemma 2.3(1) and the nesting property in the removal process reduces to the case that  $a(\Delta) \cong \nu^{-1}a(\Delta')$ . Now, showing not satisfying non-overlapping property is simply a reformulation of locally minimizability by Lemma 6.2.

Suppose  $\mathfrak{r}(\{\Delta \cap \Delta', \Delta \cup \Delta'\}, \mathfrak{h}) = \mathfrak{r}(\{\Delta, \Delta'\}, \mathfrak{h})$ . By Lemma 2.3(1), it again reduces to  $a(\Delta) \cong \nu^{-1}a(\Delta')$ . It then follows from Lemma 6.1 that  $(\{\Delta, \Delta'\}, \mathfrak{h})$  is locally minimizable and so this gives the non-overlapping property.  $\Box$ 

### 9.2. Intermediate segment property.

**Definition 9.4.** Let  $\mathfrak{h} \in \operatorname{Mult}_{\rho}$ . Let  $\Delta \in \operatorname{Seg}_{\rho}$  be admissible to  $\mathfrak{h}$ . Let  $\Delta' \in \operatorname{Seg}_{\rho}$  linked to  $\Delta$  with  $\Delta' > \Delta$ . We say that the triple  $(\Delta, \Delta', \mathfrak{h})$  satisfies the *intermediate segment* property if there exists a segment  $\widetilde{\Delta}$  in  $\mathfrak{h}$  such that

(9.3) 
$$a(\Delta) \le a(\widetilde{\Delta}) < a(\Delta'), \text{ and } b(\Delta) \le b(\widetilde{\Delta}) < b(\Delta').$$

9.3. Criteria in terms of  $\eta$ -invariants. Let  $\mathfrak{h} \in \text{Mult}_{\rho}$ . For a segment  $\Delta = [a, b]_{\rho}$  admissible to  $\mathfrak{h}$ , note that, by Theorem 2.5,  $\varepsilon_{\Delta}(\mathfrak{hd}(\pi)) = \varepsilon_{\Delta}(\pi)$ .) Let

(9.4) 
$$\eta_{\Delta}(\mathfrak{h}) = (\varepsilon_{[a,b]_{\rho}}(\mathfrak{h}), \varepsilon_{[a+1,b]_{\rho}}(\mathfrak{h}), \dots, \varepsilon_{[b,b]_{\rho}}(\mathfrak{h})).$$

The  $\eta$ -invariant defined above plays an important role in defining a notion of generalized GGP relevant pairs in [Ch22+b].

**Proposition 9.5.** Let  $\mathfrak{h} \in \text{Mult}_{\rho}$ . Let  $\Delta \in \text{Seg}_{\rho}$  be admissible to  $\mathfrak{h}$ . Let  $\Delta' \in \text{Seg}_{\rho}$  be linked to  $\Delta$  with  $\Delta' > \Delta$ . Then the following conditions are equivalent:

- (1) The triple  $(\Delta, \Delta', \mathfrak{h})$  satisfies the non-overlapping property.
- (2)  $\eta_{\Delta'}(\mathfrak{h}) = \eta_{\Delta'}(\mathfrak{r}(\Delta, \mathfrak{h})).$
- (3) The triple  $(\Delta, \Delta', \mathfrak{h})$  satisfies the intermediate segment property.

*Proof.* We first prove (3) implies (2). Suppose (3) holds. We denote by

 $\Delta_1, \ldots, \Delta_r$ 

the removal sequence for  $(\Delta, \mathfrak{h})$ . Using (3) and (4) of the removal process in Definition 2.1, those  $\Delta_1, \ldots, \Delta_r$  in  $\mathfrak{h}$  are replaced by their respective truncations, denoted by

$$\Delta_1^{tr},\ldots,\Delta_r^{tr}.$$

By using the intermediate segment property and the minimality condition in the removal process, there exists a segment of the form (9.3) in the removal sequence for  $(\Delta, \mathfrak{h})$ . Let  $i^*$  be the smallest index such that  $\Delta_{i^*}$  satisfies (\*). Note that, by considering  $a(\Delta_i)$ ,

$$\Delta_1,\ldots,\Delta_{i^*-1}$$

do not contribute to  $\eta_{\Delta'}(\mathfrak{h})$  by definitions. By the definition of truncation and (9.3) for  $\Delta_{i^*}$ , we have that  $\Delta_1^{tr}, \ldots, \Delta_{i^*-1}^{tr}$  also do not contribute to  $\eta_{\Delta'}(\mathfrak{r}(\Delta, \mathfrak{h}))$ . From the choice of  $\Delta_{i^*}$  and the nesting property, we also have that, by considering  $b(\Delta_i)$ ,

$$\Delta_{i^*},\ldots,\Delta_r$$

do not contribute to  $\eta_{\Delta'}(\mathfrak{h})$ , and similarly,  $\Delta_{i^*}^{tr}, \ldots, \Delta_r^{tr}$  do not contribute to  $\eta_{\Delta'}(\mathfrak{r}(\Delta, \mathfrak{h}))$ . Thus, we have that

$$\eta_{\Delta'}(\mathfrak{h}) = \eta_{\Delta'}(\mathfrak{r}(\Delta, \mathfrak{h}))$$

We now prove (2) implies (1). Again write  $\Delta' = [a', b']_{\rho}$ . Suppose  $(\Delta, \Delta', \mathfrak{h})$  does not satisfy the nonoverlapping property. Again, denote by

$$\Delta_1,\ldots,\Delta_r$$

the removal sequence for  $(\Delta, \mathfrak{h})$ . Let  $\Delta_l$  be the shortest segment in the removal sequence containing  $\nu^{-1}a(\Delta')$ . Note that

$$\Delta_1, \ldots, \Delta_{l-1}, \Delta_l$$

do not contribute to  $\eta_{\Delta'}(\mathfrak{h})$  (by considering  $a(\Delta_i)$ ) and similarly,

$$\Delta_1^{tr}, \ldots, \Delta_{l-1}^{tr}$$

do not contribute to  $\eta_{\Delta'}(\mathfrak{r}(\Delta,\mathfrak{h}))$ . However,  $\Delta_l^{tr}$  contributes to  $\eta_{\Delta'}(\mathfrak{r}(\Delta,\mathfrak{h}))$ . This causes a difference of 1 in the coordinate  $\varepsilon_{\Delta'}$  for  $\eta_{\Delta'}(\mathfrak{h})$  and  $\eta_{\Delta'}(\mathfrak{r}(\Delta,\mathfrak{h}))$ .

It remains to see the following claim:

Claim: For k > l,  $\Delta_k$  contributes to  $\eta_{\Delta'}(\mathfrak{h})$  if and only if  $\Delta_k^{tr}$  contributes to  $\eta_{\Delta'}(\mathfrak{r}(\Delta,\mathfrak{h}))$ .

Proof of claim: If  $\Delta_k$  does not contribute to  $\eta_{\Delta'}(\mathfrak{h})$ , then  $b(\Delta_k) < b(\Delta')$  and so  $b(\Delta_k^{tr}) < b(\Delta')$  (or  $\Delta_k^{tr}$  is dropped or a empty set). This implies that  $\Delta_k^{tr}$  does not contribute to  $\eta_{\Delta'}(\mathfrak{r}(\Delta,\mathfrak{h}))$ .

On the other hand, if  $\Delta_k$  contributes to  $\eta_{\Delta'}(\mathfrak{h})$ , then  $b(\Delta_k) \geq b(\Delta')$ . Note that  $\Delta_k^{tr}$  is non-empty by using  $\Delta < \Delta'$ . Thus we also have  $b(\Delta_k^{tr}) \geq b(\Delta')$ . We also have that  $\Delta_k^{tr}$ contributes to  $\eta_{\Delta'}(\mathfrak{r}(\Delta,\mathfrak{h}))$ . This completes proving the claim.

Note that  $(1) \Rightarrow (3)$  follows from the segment involved in the definition of overlapping property. Thus, we also have  $(1) \Rightarrow (2)$ .

We now consider (3)  $\Rightarrow$  (1). Among those segments in  $\mathfrak{h}$  satisfying (9.3), we pick the  $\prec^{L}$ minimal one  $\widetilde{\Delta}^*$  (see Section 2.3 for  $\prec^{L}$ ). Note that such segment also satisfies  $\nu^{-1}a(\Delta') \in \widetilde{\Delta}^*$  and  $\Delta' \not\subset \widetilde{\Delta}^*$ . Now (3) implies that at least one segment in the removal sequence for  $(\Delta, \mathfrak{h})$  contains a segment of the form (9.3), and so one uses the nesting property in the removal sequence to show the non-overlapping property.

**Example 9.6.** Let  $\mathfrak{h} = \{[0, 5]_{\rho}, [3, 8]_{\rho}\}.$ 

- Let  $\Delta = [0,3]_{\rho}$  and let  $\Delta' = [3,6]_{\rho}$ . In such case,  $\eta_{\Delta'}(\mathfrak{h}) = (1,0,0,0)$ . Similarly,  $\eta_{\Delta'}(\mathfrak{r}(\Delta,\mathfrak{h})) = \eta_{\Delta'}(\{[3,8]_{\rho},[4,5]_{\rho}\}) = (1,0,0,0)$ .
- Let  $\Delta = [0,3]_{\rho}$  and let  $\Delta' = [3,4]_{\rho}$ . In such case,  $\eta_{\Delta'}(\mathfrak{h}) = (1,0)$ . And  $\eta_{\Delta'}(\mathfrak{r}(\Delta,\mathfrak{h})) = \eta_{\Delta'}(\{[3,8]_{\rho},[4,5]_{\rho}\}) = (1,1)$ .

A consequence of Proposition 9.5 is the following:

**Corollary 9.7.** Suppose the triple  $(\Delta, \Delta', \mathfrak{h})$  satisfies the non-overlapping property. Write  $\Delta' = [a', b']_{\rho}$ . Then, for any segment  $\widetilde{\Delta}$  linked to  $\Delta$  and of the form  $[\widetilde{a}, b']_{\rho}$  for  $\widetilde{a} \geq a'$ , the triple  $(\Delta, \widetilde{\Delta}, \mathfrak{h})$  also satisfies the non-overlapping property.

### References

- [AGRS10] A. Aizenbud, D. Gourevitch, S. Rallis and G. Schiffmann, Multiplicity one theorems, Ann. of Math. (2) 172 (2010), no. 2, 1407-1434.
- [BZ76] I. N. Bernstein and A. V. Zelevinsky, Representations of the group GL(n, F), where F is a nonarchimedean local field, Russian Math. Surveys 31:3 (1976), 1-68.
- [BZ77] I. N. Bernstein and A. V. Zelevinsky, Induced representations of reductive p-adic groups, I, Ann. Sci. Ecole Norm. Sup. 10 (1977), 441-472.
- [Ch21] K.Y. Chan, Homological branching law for (GL<sub>n+1</sub>(F), GL<sub>n</sub>(F)): projectivity and indecomposability, Invent. math. (2021). https://doi.org/10.1007/s00222-021-01033-5
- [Ch24] K.Y. Chan, On the product functor on inner forms of the general linear group over a non-Archimedean local field, Transformation Groups (2024).
- [Ch25] K.Y. Chan, Construction of Bernstein-Zelevinsky derivatives and highest derivative multisegments I: Reduction to combinatorics, Trans. Amer. Math. Soc. Ser. B 12 (2025), 851-909.
- [Ch22+b] K.Y. Chan, Quotient branching law for *p*-adic ( $GL_{n+1}, GL_n$ ) I: generalized Gan-Gross-Prasad relevant pairs, arXiv:2212.05919 (2022)
- [Ch22+c] K.Y. Chan, Duality for the Gan-Gross-Prasad relevant pairs for *p*-adic GL, arXiv:2210.17249 (2022).
- [Ch24+e] K.Y. Chan, Construction of Bernstein-Zelevinsky derivatives and highest derivative multisegments III: properties of minimal sequences, preprint (2024).
- [CW25] Kei Yuen Chan and Daniel Kayue Wong, On the Lefeschetz principle for  $GL(n, \mathbb{C})$  and  $GL(m, \mathbb{Q}_p)$ . to appear in Israel Journal of Mathematics.
- [De23] Deng, T. Study of multiplicities in induced representations of through a symmetric reduction. manuscripta math. 171, 23â€"72 (2023). doi.org/10.1007/s00229-022-01375-1
- [Ja07] C. Jantzen, Jacquet modules of p-adic general linear groups, Represent. Theory 11 (2007), 45-83 https://doi.org/10/1090/S1088-4165-0700316-0
- [LM14] E. Lapid and A. Mínguez. On a determinantal formula of Tadić. Amer. J. Math. 136 (2014): 111-142.
- [LM16] E. Lapid, A. Mínguez, On parabolic induction on inner forms of the general linear group over a non-Archimedean local field, Sel. Math. New Ser. (2016) 22, 2347-2400.
- [LM22] E. Lapid, Mínguez, A binary operation on irreducible components of Lusztig's nilpotent varieties II: applications and conjectures for representations of GLn over a non-archimedean local field, to appear in Pure and Applied Mathematics Quarterly.
- [HM08] Marcela Hanzer, Goran Muić, On an algebraic approach to the Zelevinsky classification for classical p-adic groups, Journal of Algebra, Volume 320, Issue 8, 2008, Pages 3206-3231, doi.org/10.1016/j.jalgebra.2008.07.002.
- [Ma13] Matić, I., Jacquet modules of strongly positive representations of the metaplectic group Sp(n), Trans. Amer. Math. Soc. 365 (2013), 2755-2778. doi.org/10.1090/S0002-9947-2012-05725-4
- [MT15] Matić, I., Tadić, M. On Jacquet modules of representations of segment type. manuscripta math. 147, 437–476 (2015). https://doi.org/10.1007/s00229-015-0727-9
- [MW86] C. Mœglin, J.-L. Waldspurger, Sur l'involution de Zelevinski. J. Reine Angew. Math. 372, 136-177 (1986), doi.org/10.1515/crll.1986.372.136
- [Mi09] A. Mínguez, Surl'irréducitibilité d'une induite parabolique, 2009, no. 629, Crelle's Journal, doi.org/10.1515/CRELLE.2009.028
- [St12] R. P. Stanley, Enumerative combinatorics, vol. 1, Cambridge Studies in Advanced Mathematics 49 (Cambridge University Press, Cambridge, 2012). doi.org/10.1007/978-1-4615-9763-6
- [Ze80] A. Zelevinsky, Induced representations of reductive p-adic groups II, Ann. Sci. Ecole Norm. Sup. 13 (1980), 154-210.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG *Email address*: kychan10hku.hk